



# An Overview of the Importance of Biochar in Sustainable Agriculture

Emili Adak <sup>a</sup>, Sonai Halder <sup>a</sup>, Biresk Koley <sup>a</sup>,  
Soumyajit Biswas <sup>a</sup>, Sudip Sengupta <sup>a\*</sup>,  
Suprabuddha Kundu <sup>a</sup> and Tanmoy Sarkar <sup>a</sup>

<sup>a</sup> School of Agriculture, Swami Vivekananda University, Barrackpore, 700121, West Bengal, India.

## Authors' contributions

*This work was carried out in collaboration among all authors. Author EA designed the study, and wrote the first draft of the manuscript. Authors SH, BK, and SB performed the data collection, and literature searches. Author SS managed the entire exercise and revised the final version of the manuscript. Authors SK and TS supervised the work. All authors read and approved the final manuscript.*

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i6956>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117100>

Review Article

Received: 23/03/2024

Accepted: 27/05/2024

Published: 02/06/2024

## ABSTRACT

Sustainable agriculture is essential for addressing the global food security challenges, environmental degradation, and climate change. Biochar, a carbon-rich material produced through the pyrolysis, has gained significant attention as a promising tool to enhance sustainability of agricultural systems. Biochar serves as a valuable soil amendment by improving soil structure and enhancing its water-holding capacity, nutrient retention, and microbial activity. These improvements translate into increased soil fertility and reduced nutrient runoff, mitigating the risk of water pollution. Furthermore, biochar's porous structure acts as a reservoir for beneficial microorganisms and can potentially contribute to the suppression of soil-borne pathogens. In the context of climate change, biochar plays a dual role. Firstly, it acts as a carbon sink, sequestering carbon in the soil for hundreds to thousands of years, thus mitigating greenhouse gas emissions. Secondly, biochar application can lead to reduced emissions of nitrous oxide, a potent greenhouse gas associated

\*Corresponding author: E-mail: [sudips@svu.ac.in](mailto:sudips@svu.ac.in), [sngpta.sudip1993@gmail.com](mailto:sngpta.sudip1993@gmail.com);

**Cite as:** Adak, E., Halder, S., Koley, B., Biswas, S., Sengupta, S., Kundu, S., & Sarkar, T. (2024). An Overview of the Importance of Biochar in Sustainable Agriculture. *Journal of Advances in Biology & Biotechnology*, 27(6), 924–937. <https://doi.org/10.9734/jabb/2024/v27i6956>

with synthetic fertilizer use. Crop productivity and yield stability are critical factors in sustainable agriculture. Biochar application has been shown to enhance plant growth by improving nutrient availability, root development, and stress tolerance. Additionally, it can increase crop resilience to extreme weather events, helping farmers adapt to changing climatic conditions. However, the successful integration of biochar into agricultural systems requires careful consideration of feedstock selection, pyrolysis processes, and application methods, as well as site-specific factors. Nonetheless, continued research, development, and knowledge dissemination are essential to maximize positive impacts of biochar in diverse agricultural contexts. The current review article aims at understanding the production and applications of biochar for sustainability of agricultural production systems and minimizes the carbon emissions through sequestration.

**Keywords:** Biochar; sustainability; soil fertility; C sequestration; productivity.

## 1. INTRODUCTION

The world's population is ever increasing posing challenges to the global agricultural system. In order to feed the expanding population and meet the steadily rising demand for food, the farming system need to be more dependent on technology and chemical inputs [1]. Over the years, agricultural soil has seen an increase in the amount of fertilizer and pesticide application to increase crop yields [2]. During prolonged application, leaching losses of pesticides and fertilizers may occur, reducing soil fertility and polluting the environment. In addition to reducing soil fertility and increasing farming costs, nutrient loss from agricultural soils can also increase soil acidification and reduce crop yields [3]. In lieu of commercially prepared items, the idea on natural residue and organic substances has focused the attention of the scientific community and farmers [4]. One of the results of scientific research is biochar, which is crucial for establishing sustainable agriculture and the environment [5].

An effective way to sequester carbon to combat climate change and global warming is through the use of biochar. When incorporated into the soil, it is extremely long-lasting and can persist there for hundreds to thousands of years [6]. Additionally, it is being investigated for environmental rehabilitation, reducing pollutant mobility in polluted soils, and lessening the change of dangerous elements to agronomic crops [7]. The physical, chemical, and biological qualities of the soil are enhanced by the addition of biochar, which increases the amount of organic matter in the soil. In order to change the agricultural environment, biochar can either positively or negatively impact the microbial growth in the soil [8].

The Terra Preta, commonly referred to as "Indian black earth," was the first instance of biochar

being used to improve soil. Its black hue, high aggregate stability due to the presence of more carbon, and high nutritional content linked to an increase in microbial presence make it easy to identify [9]. The Terra Preta has been discovered to have higher levels of nutrients as well as superior soil rigidity and structure, which results in a more secure organization of soil particles [10]. Flakes from several varieties of mica were found in the soil sub-layers of the assessed Terra Preta [11]. There are several types of acidobacteria, and the Terra Preta has a 25% greater diversity of bacterial species than other soils [12].

## 2. BIOCHAR AND ITS CHARACTERISTICS

According to Tan et al. [13], biochar is a carbon-rich solid produced by heating biomass with minimal or no oxygen. It has a functional group, a porous carbonaceous structure, and an aromatic surface. An efficient waste management solution can be obtained from various origins, including agricultural, kitchen, forest-related, industry-generated, or animal-related waste. The techniques involved in production encompass torrefaction, gasification, pyrolysis, and hydrothermal/flash carbonization [14]. Among these methods, pyrolysis stands out as the most widely embraced option due to its straightforwardness, substantial output, and economical nature [15]. Biochar is typically made from waste residues including animal manures, agricultural wastes, and forestry wastes. These feedstocks are important because they have the ability to turn waste into biochar, which is a useful and valuable product. The usual pyrolysis temperature is between 200 to 800°C. At 300°C, *Lantana camara* was used to make biochar that was high in accessible P (0.64 mg/kg), Ca (5880 mg/kg), Mg (1010 mg/kg), and Na (1145 mg/kg) [16]. A large amount of N ranging from 23 to 635 mg/kg and P ranging from 46 to 1664 mg/kg can

be released from freshly made biochar, which is a rich source of readily available nutrients. After a day, biochar made from mallee wood was easily drainable with double-distilled water (15-20% Ca, 10-60% P, and 2% N) [17]. These examples show that biochar may have an impact on soil nutrient levels.

Biochar's adsorption capabilities are influenced by its physical and chemical characteristics. Biochar has extremely high specific surface area, oxygen-containing functional group content, and stability. The major influences on the physicochemical properties of biochar are the feedstock and pyrolysis temperature.

### 2.1 Surface Area

Biochar's specific surface area is crucial since it aids in the adsorption of chemicals. When the pyrolysis temperature was raised from 250 to 600°C, the surface area of sugarcane bagasse biochar raised from 0.56 to 14.1 m<sup>2</sup>/g. The release of volatiles from the biochar increases with the rise in pyrolysis temperature, which could be one explanation [18]. Additionally, the feedstock used affects the biochar's surface area. The biochar's specific surface area and pore structure can be improved by increasing the creation of vascular bundle structures as a result of the release of volatile matter, which is mostly created from celluloses and hemicelluloses during pyrolysis [19].

### 2.2 Surface Functional Group

The amount of hydrogen and oxygen, as well as the molar hydrogen to carbon (H/C) ratio, decrease as the pyrolysis temperature rises. High temperatures also significantly boost the removal efficiency of polar functional groups (-OH and C-O). As the pyrolysis temperature rises, the molar oxygen to carbon (O/C) ratio declines, resulting in a greater aromatic and lower hydrophilic surface [20].

### 2.3 Cation Exchange Capacity

Low nutrient loss from soils can be achieved with high biochar CEC. When the pyrolysis temperature was raised from 200 to 550°C, the CEC of biochar made from cordgrass appeared to grow from 8.1 to 44.5 cmol(p+)/kg and then fall to 32.4 cmol(p+)/kg [21]. The CEC of the sugarcane bagasse biochar increased from 6.40 cmol(p+)/kg at 250°C to 9.66 cmol(p+)/kg at 500°C before decreasing to 4.19 cmol(p+)/kg at 600°C [22].

### 2.4 pH Value

Applying biochar can increase the pH of the soil due to the pH of the material itself and by enhancing the retention of cations in the soil. Because alkali salts are released from the organic matrix of the feedstock as the temperature of the biochar is increased, the pH of the resulting material is higher [23]. For instance, the pH level of biochar generated from maize straw increased from 9.37 to 11.32 when the pyrolysis temperature was elevated from 300 to 600°C [24]. Due to their tendency to be more alkaline, biochar made from agricultural wastes aids in raising the pH of soils. Contrarily, because of the functional groups that animal wastes give, biochars made from materials like chicken litter or cow manure are much more acidic.

### 2.5 Biochar Stability


Increasing pyrolysis temperatures will increase biochar's stability. For instance, the stability of sugarcane bagasse biochar was significantly increased by increasing the pyrolysis temperature from 350 to 550°C. Additionally, biochar made from chicken manure had less stability than biochar made from sugarcane [25]. Biochar created at low temperatures can often be easily degraded, however biochar produced at high temperatures is resistant to degrading.

### 2.6 Water Holding Capacity

When biochar and fertilizer were applied together; there was a 14.6% increase in water content, demonstrating the importance of this porosity in increasing soil water capacity [26]. An average increase in WHC of roughly 18% results from the addition of biochar. It was interesting to note that 9.52% weight of sunflower husk biochar resulted in a 30% improvement in water retention capacity [27].

### 2.7 Electrical Conductivity

Biochar's EC can range greatly between 0.04 and 54.2 dSm<sup>-1</sup>. The reaction of biochar with water, which releases soluble chemicals, is thought to be the cause of the rise in EC [28]. According to studies, adding biochar causes the electrical conductivity of acidic red colored soil to rise. Reduced EC in saline soils may result from the physical entrapment of salts inside the pores of biochar [29].



**Fig. 1. Benefits of the use of biochar in soil-plant systems**

The pH of the soil was raised and dissolved organic matter was released when biochar made from cacao shell and rice husk was heated to 600 and 500°C, respectively [30]. Humic and fulvic acids' carbon content increased when 10% of biochar made from cow and poultry manure was added to a composting mixture. Corn straw and mushroom trash can be added to charcoal to hasten the biodegradation of polycyclic aromatic hydrocarbons [22]. The richness in organic carbon and alkaline nature of biochar also improve CEC, which increases the capacity to adsorb heavy metals. Biochar could increase porosity by 14 to 64% while reducing soil bulk density by 3 to 31%. Ma et al. [31] reported a 35% increase in the MWD following the application of 8 Mg ha<sup>-1</sup> biochar derived from corn straw and peanut hulls in China's clay loam soil. Due to its macropores and reduced surface area, it demonstrates a promising behavior of water and nutrient holding capacity in sandy soil [32]. Biochar therefore has multifaceted role in agricultural systems (Fig 1).

### 3. EFFECT OF BIOCHAR ON SOIL MICROBES

The kind of soil and crop has an impact on how biochar affects the activity of soil bacteria. In comparison to untreated soils, the diversity of the soil bacterial population increased by 25% after biochar amendment. The application of rice straw biochar greatly reduced the Actinobacteria and Ascomycota fungal communities [33]. When combined with PGPR, rice husk biochar applied at a rate of 3.6 g/kg soil was able to considerably boost rice plant output. Phosphorus and nitrogen uptake by maize plants were increased by 61.5% and 23.1%, respectively, by biochar in conjunction with *Lysinibacillus fusiformis* [34]. Rice plants' susceptibility to root knot nematode and *Fusarium* chlamydospore infection in tomato plants were both significantly reduced by biochar. With the help of defence-related gene expression, the biochar amendment may result in systemic resistance to fungi [35].

Increased levels of phenol oxidase and  $\beta$ -glucosidase were seen in mangrove sediments after biochar additions, while levels of peroxidase, N-acetyl-glucosaminidase, and acid phosphatase were decreased. Acid phosphatase (32%), alkaline phosphatase (22.8%), and fluorescein hydrolases (50%) showed the greatest increases in soil enzyme activity when *Eichornia* biochar was applied at a concentration of 20 g/kg soil [36].

Research initiatives revealed that 88.6% of L-cyfluthrin was degraded within 5 days by *Brevibacterium aureum* DG-12, a novel bacterial strain discovered in active sludge that can breakdown and use cyfluthrin as a growth substrate.  $\alpha$ -endosulfan,  $\beta$ -endosulfan, and endosulfan sulfate were all digested to 94.1%, 84.5%, and 80.1%, respectively by *Achromobacter xylosoxidans* strain C8B. Microbial consortia may be an environmentally benign technique for pesticide breakdown [37].

#### 4. INTEGRATING BIOCHAR WITH SOIL FERTILITY

The incorporation of biochar improves soil fertility (Table 1) by facilitating the biological cycle of phosphorus and nitrogen. The nitrogen fixation rose from 50% to 72% with the addition of 90 g/kg of biochar [38]. It appears that nitrogen, in the form of  $\text{NH}_3$ , is less likely to leach when using biochar. The addition of biochar to soil has significant effects on the ammonium ( $\text{NH}_4^+$ ) concentration through various processes, including  $\text{NH}_4^+$  assimilation and conversion to nitrate ( $\text{NO}_3^-$ ). Biochar effectively absorbs  $\text{NH}_4^+$  through oxygenated carbonyl and carboxyl groups, reducing its availability for nitrification and leading to higher soil  $\text{NO}_3^-$  levels [39]. Applying wheat straw biochar at rates of 0.5% to 1% was sufficient to stop various types of N leaching. It may be simple to increase the organic matter content of soils by using biochar [40]. For crops unable to fix their own nitrogen, biochar also increases the amount of nitrogen available for plant absorption. Biochar improved soil organic matter's physical security by enhancing C storage in macroaggregates of the fine-coarse soil. Sandalwood biochars from *Eucalyptus saligna* improved the mineralization of native organic C, but not in soil with a clayey texture [41]. Biochar works as a retaining agent for P and stops P from sandy soil from leaching or running off. Due to the substantial soil nutrient absorption by biochar, it acts as a slow-release fertilizer [42]. If the soil has a coarse texture and

little organic matter, biochar is a viable amendment for stabilizing aggregates. When forming aggregates and preventing soil deterioration, biochar may operate as a binding agent for organic materials. The slow oxidation properties of biochar provide the basis for the long-term effects on soil aggregation [43]. The majority of the time, plants can easily absorb the potassium present in biochar from its original source. Biochar application not only enhances soil K availability but also stimulates the proliferation of potassium-dissolving bacteria in Alfisols and Entisols. This, in turn, results in increased K availability, reducing the need for additional mineral K fertilizers [44]. The number of cyanobacteria associated with nitrogen fixation & saccharibacteria engaged in organic compound breakdown increased due to the usage of biochar-based fertilizers, thereby improving soil fertility. The result has positive influence in crop yield and production as well (Table 2).

#### 5. EFFECT OF BIOCHAR ON SOIL QUALITY

Soil quality plays a significant role in improving the status of crop.  $\text{Na}^+$  ions reduce the salinity of the soil; biochar binds to the  $\text{Na}^+$  ions and prevents the plant from absorbing them [35]. In order to increase the development of maize, the excessive salinity of central Chinese soils was reduced using a mixture of wheat straw biochar and poultry manure [55]. The use of maize stalk biochar increased partial factor production and agronomic efficiency for nitrogen in calcareous sandy soils with zucchini plants, but had no discernible impacts on phosphorus. The amount of ammonia-oxidizing bacteria in acidic soils was decreased by biochar. In acidic red soils, adding 10 g/kg of rice straw biochar raised the pH, P availability, and  $\text{Al}^{3+}$  reduced exchangeable [56].


When biochar was added to soil, the impacts of a higher pH, dispersed metal ions that were liberated, and the active groups on mineral surfaces all worked together to catalyse the hydrolysis of pesticides. According to Zhang et al. [57], biochar generated from pig manure can hydrolyze carbaryl and atrazine at 350 and 700°C with a result of 59.1% and 90.6%, respectively. Due to the catalytic properties of biochar, pesticide hydrolysis may be enhanced; however, the increased pesticide sorption may impede this improvement. When pig dung biochar produced at 350 °C after 40 days of incubation was applied at a rate of 0.5%, the

**Table 1. Nutrient retention by biochar and their effect of feedstock, pyrolysis temperature and application rate**

| Nutrient                      | Feedstock                 | Pyrolysis Temperature (°C)                                    | Rate of Biochar Application                                   | Adsorption Capacity (mg/kg) | References |
|-------------------------------|---------------------------|---|---|-----------------------------|------------|
| NH <sub>4</sub> <sup>+</sup>  | Brazilian pepperwood      | 300   | 0.1g 50m/L  | 190                         | [45]       |
|                               |                           | 450   |   | 785                         |            |
|                               |                           | 600   |   | 595                         |            |
|                               | <i>Arundo donax</i> L.    | 300   | 0.35g 12m/L   | 2101.9                      | [46]       |
|                               |                           | 350   |   | 1432.6                      |            |
|                               |                           | 400   |   | 1043.4                      |            |
| NO <sub>3</sub> <sup>-</sup>  | Brazilian pepperwood      | 400   | 0.8g 12m/L  | 362.8                       | [45]       |
|                               |                           | 500   |   | 371.8                       |            |
|                               |                           | 600   |   | 20.6                        |            |
|                               | <i>Arundo donax</i> L.    | 500   | 0.3g 8m/L   | 171.8                       | [46]       |
|                               |                           | 600   |   | 533.5                       |            |
|                               |                           | 600   |   | 333                         |            |
| PO <sub>4</sub> <sup>3-</sup> | Wheat straw               | 350-550   | 1%  | 625                         | [47]       |
|                               |                           |   | 5%  | 769                         |            |
|                               |                           |   | 10%   | 477.4                       |            |
| K <sup>+</sup>                | Sugarcane bagasse         | 450   | 0.1g 50m/L  | 19.5                        | [45]       |
|                               |                           | 300   |   | 20.3                        |            |
|                               |                           | 350   |   | 12.5                        |            |
|                               | <i>Spartina spartinae</i> | 550   | A solution flow rate of 0.30–0.35 mL/min; 50–60 mg of biochar | 6.6                         | [21]       |
|                               |                           | 300   |   | 10.9                        |            |
|                               |                           | 350   |   | 1.2                         |            |
| <i>Pinus taeda</i>            | 650                       | A solution flow rate of 0.30–0.35 mL/min; 50–60 mg of biochar |   |                             |            |
|                               |                           |   |   |                             |            |
|                               |                           |   |   |                             |            |

**Table 2. Dynamics of biochar and its effects on crop**

| SI No | Type of Biochar               | Temperature | Area of Study | Rate of Application   | Soil Type                           | Effect on Crop Yield                               | References |
|-------|-------------------------------|-------------|---------------|-----------------------|-------------------------------------|--|------------|
| 1     | Wheat Straw                   | 300–500°C   | China         | 3% w/w                | Psammaquent and Plinthudult         | Rice   | [48]       |
| 2     | Rice Straw and Corn Stalk     | 450°C       | China         | 1, 2 & 4 t/ha         | Inceptisol                          | Corn, Peanut Sweet potato                          | [49]       |
| 3     | Cow Manure                    | 600°C       | Japan         | 0, 10, 15 and 20 t/ha | Sandy soil                          | Maize  | [50]       |
| 4     | Rice Husks                    | 450 °C      | China         | 0, 10, 25 and 50 t/ha | Upland soil and paddy soil          | Rice, Wheat  | [51]       |
| 5     | Maize Stover                  | 600 °C      | USA           | 0, 1, 3, 12 & 30 t/ha | Kendaia silt loam                   | NA   | [52]       |
| 6     | Woodchips                     | 700°C       | USA           | 5% w/w                | Clay texture and sandy loam texture | NA   | [53]       |
| 7     | <i>Eucalyptus polybractea</i> | 550°C       | UK            | 10 t/ha               | Ferrosol Soil                       | Cauliflower, peas, Broccoli- no significant effect | [54]       |



**Fig. 2. Reaction of biochar in retention of nutrients and heavy metals**

breakdown efficiency of carbaryl increased from 55.0% to 75.0% in the unsterile soil. By affecting the sorption and desorption processes, biochar decreased the microbial atrazine mineralization, hence lowering the atrazine's bioavailability. In soil that has been altered with biochar, pesticide mobility, volatilization, leaching, and uptake by plants can all be reduced. The characteristics of biochar, including its level of organic carbon, fragrance, specific surface area, and ash content, have a significant impact on its capacity to bind pesticides [58].

Metals in a soil are immobilized by biochar because of its lower particle size. This immobilization is brought on by the biochar's increased surface area, which finally results in less metal absorption by plant tissues. When applied at 5%, tobacco stalk biochar decreased the uptake of Cd and Zn by 64.2 and 94.9%, respectively [59]. Rice grain methyl mercury content was decreased by applying biochar made from rice straw and heated to 600 °C at concentrations of 1% to 4% by 49-92%. Rice plants took up less As when Mn modified biochar was applied to As-contaminated soil, and the amount of amino acids rose by 9% [60]. Using biochar in conjunction with other chemicals, including limestone, decreased the uptake of Cd in wheat and rice plant straw and grain by more than 80%. Plant uptake of metals was frequently decreased by applying biochar at depths of 0 to 15 cm. The amount of chlorpyrifos and fipronil that *Allium tuberosum* absorbed after being exposed to soil with 1% straw chip biochar was

reduced by 19 and 48%, respectively [61]. Chinese cabbage plants' uptake of PAHs was reduced when wheat straw biochar was added at a rate of 2%. The absorption of nanoparticles of cerium oxide by zucchini crops could also be decreased by applying biochar made from pecan shells at a rate of 5% [62].

Polar gases, water-soluble metals, and different organic contaminants are normally all absorbed by biochar. Generally speaking, biochars with greater aromatic qualities and greater C condensation are more resilient. Biochar efficiently binds heavy metal cations from water and immobilizes elements of heavy metal in soil [13,63]. By interacting with the applied biochar and adhering to the pore surfaces, heavy metal ions in the soil may be changed into hydroxides, carbonates, and phosphate precipitates (Fig 2). As a result of this process, the proportion of heavy metals those are water-soluble and bioactive in soil drops, which diminish their potential for uptake and bioaccumulation by soil organisms, including plant roots [63]. By exchanging the cations that were previously connected to its surface functional units through ion exchange, biochar is able to adsorb heavy metal ions from the soil solution. Biochar sorbs Cd, Cu, Ni, and Pb mostly through surface complexation, which makes use of ligand-like surface functional elements like carboxylic, hydroxyl, and phenolic groups [64]. The addition of alkalinity from biochar often results in an increase in soil pH. Increased pH promotes heavy metal hydrolysis, which results in the



creation of metal hydroxide precipitates. This lowers the amount of metal ions that is soluble in water and helps minimize soil heavy metal contamination [65]. Organic non-ionic compounds are adsorbable on biochar via surface adsorption along with partitioning. Because of their higher surface energy, biochar micropores are where the adsorption of organic pollutants begins [64]. Metal ions can combine with polar and charged organic molecules to produce complexes that can either be deposited on the surface of the biochar or precipitate in the soil.

## 6. EFFECT OF BIOCHAR ON GREENHOUSE GASES

Biochar is also considered as a boon as it is possible to reduce CO<sub>2</sub> emissions by 1.67 tonnes of CO<sub>2</sub> equivalent every tonne of feedstock utilized during application. It is important to note that woody biomass-derived biochar delivers greater emission reductions because of its increased energy content, which can counteract GHG emissions created during the biomass burning process [66]. The concentration of soil organic carbon (SOC) is also affected by the maturation of biochar in the soil. For instance, Zhang et al. [67] found that the application of wheat straw biochar (40 Mg ha<sup>-1</sup>) in rice fields led to a 57% increase in SOC concentrations. In calcareous soil with low OC, they reported that addition of biochar at 20 and 40 Mg ha<sup>-1</sup> integrated with nitrogen fertilizers increased SOC by 25 to 42.2%. It was proposed that biochar might deliver a maximum of 15-35% of the carbon dioxide reduction target by using residue biomass and growing additional crops expressly for biochar production. Additionally, iron- or calcium-treated biochar demonstrated its value in increasing the stability of soil aggregates and lowering carbon emissions with improved carbon sequestration abilities [67]. Because biochar decomposes organic carbon more slowly than straw biomass, applying it to paddy fields minimizes CH<sub>4</sub> emissions from those areas [66]. The addition of biochar leads in less CH<sub>4</sub> being released from the carbonaceous substances than straw biomass and inhibits the spread of methanogenic bacteria by providing a comfortable environment in its porous structure.

Additionally, biochar enhances the production of N<sub>2</sub>O reductase genes in denitrifying bacteria and encourages microbial N immobilization [66]. This makes it easier for N<sub>2</sub>O to be converted to N<sub>2</sub> during denitrification, which lowers emissions of

N<sub>2</sub>O from soil. N<sub>2</sub>O emissions dropped by 50% after the addition of wood biochar formed at 350 °C and 700 °C biochar, respectively. Utilizing biochar made from pine, willow, and corn, which were all created at 550 °C, decreased N<sub>2</sub>O emissions by around 40%, 46%, and 60%, accordingly [68]. Higher-temperature biochars were more effective at lowering the release of N<sub>2</sub>O than NO emissions, and lower-temperature biochars are more effective at lowering NO emissions. Crop yields increased and N<sub>2</sub>O emissions decreased after straw-derived biochar was added to sandy-loam soils for a five-year wheat and maize crop rotation, but total GWP did not change. It is believed that biochar will have the ideal characteristics for preventing global warming and changing the climate. Numerous research support the idea that applying biochar to areas used for cultivating vegetables will significantly reduce emissions of carbon dioxide, N<sub>2</sub>O, and methane gasses [69,70].

## 7. EFFECT OF BIOCHAR IN SUSTAINABILITY

Lichtfouse et al. [71] revealed the concept of "sustainable agriculture" as the combination of socioeconomic research with biological, chemical, physical, and ecological functions in order to create new agricultural techniques that are secure and ecologically sound. By conserving and upholding all of its natural resources, such as conserving soil fertility, protecting surface and below-ground assets, creating renewable energy sources, and looking for ways to adapt farming practices to climate change, sustainable agriculture enables agro farming to sustain itself over an extended period of time [72]. Agro farming needs to take social groupings and the environment's sustainability into account.

The main challenges facing the existing agro farming methods are to increase crop yield in a way that is environmentally sound and sustainable [73]. A more sustainable approach to farming will lessen the need for organic fertilizers and maintain the productivity of agriculture. The majority of current biochar research focuses on modifying the material's characteristics to improve its ability to remove both organic as well as inorganic contaminants. Biochar exhibits amazing efficacy in decreasing contaminants including antibacterial agents, herbicides, pigments, pesticides, and heavy metals due to its outstanding surface properties, and it is essential in addressing the effects of climate change [74].

The extraordinary ability of biochar to immobilize rhizospheric heavy metals and agricultural pesticides on its vast surface and prevent their uptake by crops increases crop production [75]. It has been demonstrated that adding biochar reduces the soil's maximum hydraulic conductivity, particularly in light-textured soils [76].

## 8. CONCLUSIONS

Biochar shines as a ray of hope in our pursuit of sustainable farming, improved soil health, vigorous cultivation, and efficient climate change management. Its many advantages, including increased nutrient retention, less release of greenhouse gases, and greater soil capacity to hold water, offer a positive vision for a more robust and environmentally conscientious agricultural destiny. We can lessen our impact on our surroundings, repair damaged soils, and equip farmers to meet the hurdles of a changing climate by incorporating biochar into our farming practices. With the use of biochar, we may conserve nature without sacrificing crop output by delaying the release of resources. Biochar must be seen as more than just a fix—it must be seen as a road toward a more peaceful living with our planet as we work to fully realize its potential through continued study and creative use. We can make a big step towards sustainable farming, better ecosystems, and a more sustainable planet for future generations by embracing biochar.

## 9. LIMITATIONS OF THE USE OF BIOCHAR

There are several restrictions even if the majority of research data demonstrated the advantages of biochar treatments. Additionally, *Oryza sativa* and *Solanum lycopersicum*'s subsurface root biomass decreased in response to aged biochar [77]. It has been demonstrated that biochar has favourable soil-specific impacts. Using biochar at a high level in lentil cultivation caused the development of weeds to rise by 200% [78]. Implementation of biochar may also cause plants to blossom later. In saline and sodic soil, applying biochar and phosphorus fertilizer together may speed up the phosphate precipitation/sorption processes. This interaction may ultimately result in the plants' access to phosphorus being reduced. It has been demonstrated that the biochar source's contamination is harmful for crop development. Based on the kind, quantity, kind of soil, crop

variety, etc. of the biochar amendment, beneficial as well as detrimental impacts on crop production and fertility fluctuations might occur.

## 10. FUTURE LINE OF RESEARCH BY USE OF NANO BIOCHAR

Recently a number of studies have been done on the sustainable production of nano-biochar for land and agricultural use. A micro-sized biochar known as "dissolve" and "nano-biochar" is created during the carbonization process. It has dimensions less than or equal to a micrometer and up to a nanometer. Recent research produced nano-biochar with a diameter <5 nm. The structural modifications and physico-chemical properties of bulk biochar and nanobiochar differ from one another. The elemental compositions, aromatic/polar nature, pH, CEC, specific surface area (SSA), pore properties, and zeta potential of nano-biochar can all vary. Nano-biochar is also available in soluble and insoluble forms. Dairy manure biochar generally has a soluble carbon concentration of 14% at 100°C to about 0.2% at 700°C, with the soluble carbon content usually rising according to rising solvent pH and decreasing with rising pyrolysis temperature. As a result of rising instances of environmental degradation and decreased crop productivity caused by shifting climatic circumstances, nano-biochar becomes increasingly intriguing [79,80].

## ACKNOWLEDGEMENT

Authors are thankful to honourable Chancellor and Vice-Chancellor of Swami Vivekananda University for their constant encouragements. Authors are also thankful to honourable COO Sir for his help and supports in all means.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Mueller ND, Gerber JS, Johnston M, Ray DK, Ramankutty N, Foley JA. Closing yield gaps through nutrient and water management. *Nature*, 2012;490(7419): 254-257.
2. Pan D, Kong F, Zhang N, Ying R. Knowledge training and the change of fertilizer use intensity: Evidence from

- wheat farmers in China. *Journal of Environmental Management*. 2017;197:130-139.
3. Laird D, Fleming P, Wang B, Horton R, Karlen D. Biochar impact on nutrient leaching from a Midwestern agricultural soil. *Geoderma*, 2010;158(3-4):436-442.
  4. Ilvo P, Uog P, Slu P, Inra P, Upm P, Unitus P, Iamo P, Iea-ar P. D5.2 Participatory impact assessment of sustainability and resilience of EU farming systems. 2019: No; June 2017.
  5. Lu L, Yu W, Wang Y, Zhang K, Zhu X, Zhang Y, Wu Y, Ullah H, Xiao X, Chen B. Application of biochar-based materials in environmental remediation: From multi-level structures to specific devices. *Biochar*. 2020;2:1-31.
  6. Ayaz M, Feizienė D, Tilvikienė V, Akhtar K, Stulpinaitė U, Iqbal R. Biochar role in the sustainability of agriculture and environment. *Sustainability*, 2021;13(3):1330.
  7. Jalal F, Arif M, Akhtar K, Khan A, Naz M, Said F, Zaheer S, Hussain S, Imtiaz M, Khan MA, Wei F. Biochar integration with legume crops in summer gape synergizes nitrogen use efficiency and enhance maize yield. *Agronomy*. 2020;10(1):58.
  8. Kavitha B, Reddy PVL, Kim B, Lee SS, Pandey SK, Kim KH. Benefits and limitations of biochar amendment in agricultural soils: A review. *Journal of environmental management*. 2018;227:146-154.
  9. Glaser B, Birk JJ. State of the scientific knowledge on properties and genesis of anthropogenic dark earths in central amazonia (terra preta de Índio). *Geochimica et Cosmochimica acta*, 2012;82:39-51.
  10. Alling V, Hale SE, Martinsen V, Mulder J, Smebye A, Breedveld GD, Cornelissen G. The role of biochar in retaining nutrients in amended tropical soils. *Journal of Plant Nutrition and Soil Science*. 2014;177(5):671-680.
  11. Lima HN, Schaefer CE, Mello JW, Gilkes RJ, Ker JC. Pedogenesis and pre-colombian land use of "Terra Preta Anthrosols" ("Indian black earth") of Western Amazonia. *Geoderma*. 2002;110(1-2):1-17.
  12. Kim JS, Sparovek G, Longo RM, De Melo WJ, Crowley D. Bacterial diversity of terra preta and pristine forest soil from the Western Amazon. *Soil Biology and Biochemistry*. 2007;39(2):684-690.
  13. Tan X, Liu Y, Zeng G, Wang X, Hu X, Gu Y, Yang Z. Application of biochar for the removal of pollutants from aqueous solutions. *Chemosphere*. 2015;125:70-85.
  14. Hu F, Xu C, Ma R, Tu K, Yang J, Zhao S, Yang M, Zhang F. Biochar application driven change in soil internal forces improves aggregate stability: Based on a two-year field study. *Geoderma*. 2021;403:115276.
  15. Uday V, Harikrishnan PS, Deoli K, Zitouni F, Mahlknecht J, Kumar M. Current trends in production, morphology, and real-world environmental applications of biochar for the promotion of sustainability. *Bioresource Technology*. 2022;359:127467.
  16. Antonangelo JA, Zhang H. Heavy metal phytoavailability in a contaminated soil of northeastern Oklahoma as affected by biochar amendment. *Environmental Science and Pollution Research*. 2019;26:33582-33593.
  17. Wang K, Peng N, Lu G, Dang Z. Effects of pyrolysis temperature and holding time on physicochemical properties of swine-manure-derived biochar. *Waste and Biomass Valorization*. 2020;11:613-624.
  18. Liu X, Zhang Y, Li Z, Feng R, Zhang Y. Characterization of corncob-derived biochar and pyrolysis kinetics in comparison with corn stalk and sawdust. *Bioresource Technology*. 2014;170:76-82.
  19. Lee Y, Park J, Ryu C, Gang KS, Yang W, Park YK, Hyun S. Comparison of biochar properties from biomass residues produced by slow pyrolysis at 500 °C. *Bioresource technology*. 2013;148:196-201.
  20. Chen B, Zhou D, Zhu L. Transitional adsorption and partition of nonpolar and polar aromatic contaminants by biochars of pine needles with different pyrolytic temperatures. *Environmental Science & Technology*. 2008;42(14):5137-5143.
  21. Harvey OR, Herbert BE, Rhue RD, Kuo LJ. Metal interactions at the biochar-water interface: Energetics and structure-sorption relationships elucidated by flow adsorption microcalorimetry. *Environmental Science & Technology*. 2011;45(13):5550-5556.
  22. Ding W, Dong X, Ime IM, Gao B, Ma LQ. Pyrolytic temperatures impact lead sorption mechanisms by bagasse biochars. *Chemosphere*. 2014;105:68-74.

23. Ahmad M, Lee SS, Dou X, Mohan D, Sung JK, Yang JE, Ok YS. Effects of pyrolysis temperature on soybean stover-and peanut shell-derived biochar properties and TCE adsorption in water. *Bioresource Technology*. 2012;118:536-544.
24. Yuan JH, Xu RK, Zhang H. The forms of alkalis in the biochar produced from crop residues at different temperatures. *Bioresource Technology*. 2011;102(3): 3488-3497.
25. Cross A, Sohi SP. A method for screening the relative long-term stability of biochar. *Gcb Bioenergy*. 2013;5(2):215-220.
26. Agegnehu G, Srivastava AK, Bird MI. The role of biochar and biochar-compost in improving soil quality and crop performance: A review. *Applied Soil Ecology*. 2017;119:156-170.
27. Gluba Ł, Rafalska-Przysucha A, Szewczak K, Łukowski M, Szlązak R, Vitková J, Usowicz B. Effect of fine size-fractionated sunflower husk biochar on water retention properties of arable sandy soil. *Materials*. 2021;14(6):1335.
28. Joseph S, Cowie AL, Van Zwieten L, Bolan N, Budai A, Buss W, Lehmann J. How biochar works, and when it doesn't: A review of mechanisms controlling soil and plant responses to biochar. *Gcb Bioenergy*. 2021;13(11):1731-1764.
29. Thomas SC, Frye S, Gale N, Garmon M, Launchbury R, Machado N, Winsborough C. Biochar mitigates negative effects of salt additions on two herbaceous plant species. *Journal of Environmental Management*. 2013;129:62-68.
30. Mandal S, Pu S, Adhikari S, Ma H, Kim DH, Bai Y, Hou D. Progress and future prospects in biochar composites: Application and reflection in the soil environment. *Critical Reviews in Environmental Science and Technology*. 2021;51(3):219-271.
31. Ma N, Zhang L, Zhang Y, Yang L, Yu C, Yin G, Ma X. Biochar improves soil aggregate stability and water availability in a mollisol after three years of field application. *PLOS ONE*. 2016;11(5): e0154091.
32. Radwan NM, Marzouk ER, El-Melegy AM, Hassan MA. Improving soil properties by using biochar under drainage conditions in North Sinai. *Sinai Journal of Applied Sciences*. 2020;9(2):157-168.
33. Farrell M, Kuhn TK, Macdonald LM, Maddern TM, Murphy DV, Hall PA, Baldock JA. Microbial utilisation of biochar-derived carbon. *Science of the Total Environment*. 2013;465:288-297.
34. Rafique M, Sultan T, Ortas I, Chaudhary HJ. Enhancement of maize plant growth with inoculation of phosphate-solubilizing bacteria and biochar amendment in soil. *Soil science and plant nutrition*. 2017;63(5):460-469.
35. Akhter A, Hage-Ahmed K, Soja G, Steinkellner S. Compost and biochar alter mycorrhization, tomato root exudation, and development of *Fusarium oxysporum* f. sp. *lycopersici*. *Frontiers in Plant Science*. 2015;6:529.
36. Du Z, Wang Y, Huang J, Lu N, Liu X, Lou Y, Zhang Q. Consecutive biochar application alters soil enzyme activities in the winter wheat-growing season. *Soil Science*. 2014;179(2):75-83.
37. Singh NS, Singh DK. Biodegradation of endosulfan and endosulfan sulfate by *Achromobacter xylosoxidans* strain C8B in broth medium. *Biodegradation*. 2011;22: 845-857.
38. Rondon MA, Lehmann J, Ramírez J, Hurtado M. Biological nitrogen fixation by common beans (*Phaseolus vulgaris* L.) increases with bio-char additions. *Biology and fertility of soils*. 2007;43: 699-708.
39. Abbruzzini TF, Davies CA, Toledo FH, Cerri CEP. Dynamic biochar effects on nitrogen use efficiency, crop yield and soil nitrous oxide emissions during a tropical wheat-growing season. *Journal of Environmental Management*. 2019;252: 109638.
40. Laskosky JD, Mante AA, Zvomuya F, Amarakoon I, Leskiw L. A bioassay of long-term stockpiled salvaged soil amended with biochar, peat, and humalite. *Agrosystems, Geosciences & Environment*. 2020;3(1):e20068.
41. Jien SH, Wang CS. Effects of biochar on soil properties and erosion potential in a highly weathered soil. *Catena*. 2013;110: 225-233.
42. Dai Y, Wang W, Lu L, Yan L, Yu D. Utilization of biochar for the removal of nitrogen and phosphorus. *Journal of Cleaner Production*. 2020;257:120573.
43. Patel JS, Singh A, Singh HB, Sarma BK. Plant genotype, microbial recruitment and nutritional security. *Frontiers in Plant Science*. 2015;6:608.
44. Singh A, Singh AP, Purakayastha TJ. Characterization of biochar and their influence on microbial activities and

- potassium availability in an acid soil. Archives of Agronomy and Soil Science. 2019;65(9):1302-1315.
45. Yao Y, Gao B, Zhang M, Inyang M, Zimmerman AR. Effect of biochar amendment on sorption and leaching of nitrate, ammonium, and phosphate in a sandy soil. Chemosphere. 2012;89(11):1467-1471.
  46. Zheng H, Wang Z, Deng X, Herbert S, Xing B. Impacts of adding biochar on nitrogen retention and bioavailability in agricultural soil. Geoderma. 2013;206:32-39.
  47. Xu G, Sun J, Shao H, Chang SX. Biochar had effects on phosphorus sorption and desorption in three soils with differing acidity. Ecological Engineering. 2014;62:54-60.
  48. Muhammad N, Aziz R, Brookes PC, Xu J. Impact of wheat straw biochar on yield of rice and some properties of Psammaquent and Plinthudult. Journal of Soil Science and Plant Nutrition. 2017; 17(3):808-823.
  49. Yang Y, Ma S, Zhao Y, Jing M, Xu M, Chen J. A field experiment on enhancement of crop yield by rice straw and corn stalk-derived biochar in Northern China. Sustainability. 2015;7:13713–13725.
  50. Azeem M, Hayat R, Hussain Q, Ahmed M, Pan G, Tahir MI, Irfan M. Biochar improves soil quality and N<sub>2</sub>-fixation and reduces net ecosystem CO<sub>2</sub> exchange in a dryland legume-cereal cropping system. Soil and Tillage Research. 2019;186:172-182.
  51. Wang J, Pan X, Liu Y, Zhang X, Xiong Z. Effects of biochar amendment in two soils on greenhouse gas emissions and crop production. Plant and soil. 2012;360:287-298.
  52. Güereña D, Lehmann J, Hanley K, Enders A, Hyland C, Riha S. Nitrogen dynamics following field application of biochar in a temperate North American maize-based production system. Plant and soil. 2013;365:239-254.
  53. Lai WY, Lai CM, Ke GR, Chung RS, Chen CT, Cheng CH, Chen CC. The effects of woodchip biochar application on crop yield, carbon sequestration and greenhouse gas emissions from soils planted with rice or leaf beet. Journal of the Taiwan Institute of Chemical Engineers. 2013;44(6): 1039-1044.
  54. Boersma M, Wrobel-Tobiszewska A, Murphy L, Eyles A. Impact of biochar application on the productivity of a temperate vegetable cropping system. New Zealand Journal of Crop and Horticultural Science. 2017;45(4):277-288.
  55. Zhu Q, Wu J, Wang L, Yang G, Zhang X. Effect of biochar on heavy metal speciation of paddy soil. Water, Air, & Soil Pollution. 2014;226:1-10.
  56. Zhang P, Sun H, Yu L, Sun T. Adsorption and catalytic hydrolysis of carbaryl and atrazine on pig manure-derived biochars: Impact of structural properties of biochars. Journal of Hazardous Materials. 2013;244:217-224.
  57. Fang Q, Chen B, Lin Y, Guan Y. Aromatic and hydrophobic surfaces of wood-derived biochar enhance perchlorate adsorption via hydrogen bonding to oxygen-containing organic groups. Environmental Science & Technology. 2014;48(1):279-288.
  58. Yang X, Lu K, McGrouther K, Che L, Hu G, Wang Q, Wang H. Bioavailability of Cd and Zn in soils treated with biochars derived from tobacco stalk and dead pigs. Journal of Soils and Sediments. 2017;17:751-762.
  59. Yu Z, Qiu W, Wang F, Lei M, Wang D, Song Z. Effects of manganese oxide-modified biochar composites on arsenic speciation and accumulation in an indica rice (*Oryza sativa* L.) cultivar. Chemosphere. 2017;168:341-349.
  60. Yang XB, Ying GG, Peng PA, Wang L, Zhao JL, Zhang LJ, He HP. Influence of biochars on plant uptake and dissipation of two pesticides in an agricultural soil. Journal of Agricultural and Food Chemistry. 2010;58(13):7915-7921.
  61. Servin AD, De la Torre-Roche R, Castillo-Michel H, Pagano L, Hawthorne J, Musante C, White JC. Exposure of agricultural crops to nanoparticle CeO<sub>2</sub> in biochar-amended soil. Plant Physiology and Biochemistry. 2017;110:147-157.
  62. Ahmad M, Rajapaksha AU, Lim JE, Zhang M, Bolan N, Mohan D, Ok YS. Biochar as a sorbent for contaminant management in soil and water: a review. Chemosphere. 2014;99:19-33.
  63. Uchimiya M, Chang S, Klasson KT. Screening biochars for heavy metal retention in soil: Role of oxygen functional groups. J. Hazard. Mater. 2011;190:432–441.
  64. Ippolito JA, Berry CM, Strawn DG, Novak JM, Levine J, Harley A. Biochars reduce mine land soil bioavailable metals. J. Environ. Qual. 2017;46:411–419.
  65. Lehmann J, Cowie A, Masiello CA, Kammann C, Woolf D, Amonette JE,

- Whitman T. Biochar in climate change mitigation. *Nature Geoscience*. 2021;14(12):883-892.
66. Liu H, Xu G, Li G. The characteristics of pharmaceutical sludge-derived biochar and its application for the adsorption of tetracycline. *Science of The Total Environment*. 2020;747:141492.
67. Zhang H, Chen C, Gray EM, Boyd SE, Yang H, Zhang D. Roles of biochar in improving phosphorus availability in soils: A phosphate adsorbent and a source of available phosphorus. *Geoderma*. 2016;276:1-6.
68. Nelissen V, Saha BK, Ruyschaert G, Boeckx P. Effect of different biochar and fertilizer types on N<sub>2</sub>O and NO emissions. *Soil Biology and Biochemistry*. 2014;70:244-255.
69. Sun L, Li L, Chen Z, Wang J, Xiong Z. Combined effects of nitrogen deposition and biochar application on emissions of N<sub>2</sub>O, CO<sub>2</sub> and NH<sub>3</sub> from agricultural and forest soils. *Soil Science and Plant Nutrition*. 2014;60(2):254-265.
70. Hale SE, Nurida NL, Mulder J, Sørmo E, Silvani L, Abiven S, Cornelissen G. The effect of biochar, lime and ash on maize yield in a long-term field trial in a Ultisol in the humid tropics. *Science of the Total Environment*. 2020;719:137455.
71. Lichtfouse E, Navarrete M, Debaeke P, Souchère V, Alberola C, Ménassieu J. *Agronomy for sustainable agriculture: a review*. Sustainable Agriculture. 2009;1-7.
72. Akhtar K, Wang W, Ren G, Khan A, Enguang N, Khan A, Wang H. Straw mulching with inorganic nitrogen fertilizer reduces soil CO<sub>2</sub> and N<sub>2</sub>O emissions and improves wheat yield. *Science of the Total Environment*. 2020;741:140488.
73. Srivastav AL. Chemical fertilizers and pesticides: Role in groundwater contamination in agrochemicals detection, treatment and remediation. Butterworth-Heinemann. 2020;143-159.
74. Schiermeier Q. Putting the carbon back: The hundred billion tonne challenge. *Nature*. 2006;442(7103):620-624.
75. Bolan NS, Choppala G, Kunhikrishnan A, Park J, Naidu R. Microbial transformation of trace elements in soils in relation to bioavailability and remediation (pp. 1-56). Springer New York; 2013.
76. Brockhoff SR, Christians NE, Killorn RJ, Horton R, Davis DD. Physical and mineral-nutrition properties of sand-based turfgrass root zones amended with biochar. *Agronomy Journal*. 2010; 102(6):1627-1631.
77. Anyanwu IN, Alo MN, Onyekwere AM, Crosse JD, Nworie O, Chamba EB. Influence of biochar aged in acidic soil on ecosystem engineers and two tropical agricultural plants. *Ecotoxicology and Environmental Safety*. 2018;153:116-126.
78. Safaei Khorram M, Fatemi A, Khan MA, Kiefer R, Jafarnia, S. Potential risk of weed outbreak by increasing biochar's application rates in slow-growth legume, lentil (*Lens culinaris* Medik.). *Journal of the Science of Food and Agriculture*. 2018;98(6):2080-2088.
79. Chen T, Zhou Z, Xu S, Wang H, Lu W. Adsorption behavior comparison of trivalent and hexavalent chromium on biochar derived from municipal sludge. *Bioresource Technology*. 2015;190:388-394.
80. Rajput VD, Minkina T, Ahmed B, Singh VK, Mandzhieva S, Sushkova S, Wang B. Nano-biochar: A novel solution for sustainable agriculture and environmental remediation. *Environmental Research*. 2022;210:112891

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/117100>