



Impact of Mechanical Ventilation on Outcomes of COVID-19 Patients in Hodeidah, Yemen

Mohammed Amod Al-Kamarany ^{a,b*}, Naqib Ahmad ^b,
Abdoalrhman Almusabli ^a, Ahmed Majam ^a
and Khalid Suhail ^a

^a Center of Tropical Medicine and Epidemiology Studies, Hodeidah University (CTMES – HU), Yemen.

^b Center of Tropical Medicine and Infectious Diseases (CTMID), Al-Thawara Public Hospital Authority, Hodeidah, Yemen.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.globalpresshub.com/review-history/1612>

Original Research Article

Received: 17/04/2024

Accepted: 20/06/2024

Published: 06/08/2024

ABSTRACT

Background: Severe and critical infection of coronavirus disease 2019 (COVID -19) is characterized by Acute Respiratory Distress Syndrome (ARDS). The use of ventilators (non-invasive and mechanical), to deliver oxygen in cases of severe and critical COVID-19. Thus far, among the 10-20 % of COVID-19 patients who require treatment in an intensive care unit (ICU), about 5 % of them have been placed on mechanical ventilation (MV).

Objective: Therefore, the study aimed to estimate the impact of MV in management of COVID - 19 in Hodeidah, Yemen.

*Corresponding author: Email: alkamarany@gmail.com;

Cite as: Al-Kamarany, M. A., Ahmad, N., Almusabli, A., Majam, A., & Suhail, K. (2024). Impact of Mechanical Ventilation on Outcomes of COVID-19 Patients in Hodeidah, Yemen. *Asian Journal of Research in Medicine and Medical Science*, 6(1), 155–163. Retrieved from <https://jofmedical.com/index.php/AJRMMS/article/view/79>.

Methodology: This study was designed in a case series study that included 28 COVID-19 patients admitted to ICU, isolation department, Center of Tropical Medicine and Infectious Diseases (CTMID), AL Thawrah Public Hospital Authority, Hodeidah, Yemen as a reference hospital. 28 critical illness cases were reported in the first wave of 2020. Data included demographics, advanced life support therapies, and MV variables. The MV variables included respiratory rate (RR); fraction of inspired oxygen (FiO₂); M-V; Tidal volume (Ti); positive end-expiratory pressure (PEEP); inspiratory to expiratory ratio (I:E), flow rate; peak inspiratory pressure (PIP); plateau pressure; SaO₂; arterial pressure of oxygen (PaO₂); arterial pressure of carbon dioxide (PaCO₂), pH, and ventilator modes that were adjusted.

Results: The results of the first wave in 2020 showed 28/505 (5.54%) patients with critical illness underwent MV. Only 6/28 patients (21.4%) were recovered where a significant difference was reported ($\chi^2 = 9.143$; $p = 0.0025$) in comparing with severe cases with non-invasive ventilation namely 20/21 (95.23%) that were recovered. The main outcome for recovery was 31-day survival.

Conclusion: During the peak of the pandemic in Hodeidah, Yemen, critically ill patients with COVID-19 often required MV and mortality was very high for critical cases in the first waves, therefore the best practices of MV in COVID-19 must be improved.

Keywords: Impact; mechanical ventilator; COVID-19; Hodeidah; Yemen.

1. INTRODUCTION

“Mechanical ventilation (MV) is used to treat patients with severe coronavirus disease 2019 (COVID-19), little is known about the long-term health implications of this treatment” [1]. “Early observations suggested that COVID-19 pneumonia had a higher mortality rate than other causes of pneumonia” [2]. “There are large uncertainties with regard to the outcome of patients with COVID-19 and MV. High mortality (50-97%) was proposed by some groups, leading to considerable uncertainties with regard to outcomes of critically ill patients with COVID-19” [3].

“The mechanisms of COVID-19-associated ARDS include severe pulmonary infiltration/edema and inflammation, leading to impaired alveolar homeostasis, alteration of pulmonary physiology resulting in pulmonary fibrosis, endothelial inflammation and vascular thrombosis. Despite some distinct differences between COVID-19-associated Acute Respiratory Distress Syndrome (ARDS) and classical ARDS as defined by Berlin criteria, general treatment principles, such as lung-protective ventilation and rehabilitation concepts should be applied whenever possible. At the same time, ventilatory settings for COVID-19-associated ARDS require to be adapted in individual cases, depending on respiratory mechanics, recruitability and presentation timing” [4-8].

“An ongoing outbreak of pneumonia associated with COVID-19 started in December, 2019, in

Wuhan, China. In Hadramout, Yemen was reported in April, 2020 and in Hodeidah, Yemen was diagnosed in May 2020. Information about critically ill patients with COVID-19 infection is scarce” [9-11]. The study aimed to describe the clinical course and outcomes of critically ill patients with COVID-19 pneumonia.

2. METHODOLOGY

2.1 Study Area, Setting and Designs

The study was carried out in the intensive care unit (ICU) of the COVID-19 isolation department, Center of Tropical Medicine and Infectious Diseases (CTMID), AL-Thawrah Public Hospital Authority, Hodeidah, Yemen. The study was designed in a case series study (a retrospective study) from 1st June to 31st December 2020.

2.2 COVID-19 Confirmation

The cases were confirmed with COVID-19 using on Real Time - Polymerase Chain Reaction (RT-PCR) in the biological molecular unit of CTMID - Authority of Public Al Thawrah Hospital, Hodeidah, Yemen [12,13].

2.3 Critical Cases Diagnosis

The case definition of critical cases depended on the Yemeni national guideline that was extracted from the World Health Organization (WHO) guideline [14,15]. Different clinical investigations were used to assess the critical cases, namely radiological features, hematological features and biochemical features [16-18].

2.4 Critical Cases Management

2.4.1 Pharmacological approach

The pharmacological approach of COVID-19 critical illness was summarized in Table (1) [19].

2.4.2 Non – pharmacological approach (Mechanical ventilation)

The MV parameters for treatment of COVID-19 critical illness was summarized in Table 2 [20,21].

2.5 Data Collection and Analysis

The independent variables studied namely age, gender, clinical signs, risk factors, co-morbidity, co-infection, pharmacological medication, non – pharmacological medication (MV parameters) that were collected. Data were checked and entered in Statistical Package for Social Science (SPSS) and Microsoft Excel. The data were subsequently visualized using tables, graphs and text. Data were described through calculations, mean and percentages. Comparisons between qualitative variables were analyzed using Chi-squared test.

3. RESULTS

3.1 Sex – Age and Outcome

A total of 28 patients were confirmed undergo to MV. Males were significantly overrepresented in this group compared with females ($X^2 = 7$; $p = 0.00815$). The results showed a statistically significant of COVID-19 infection with increasing with age, where higher frequency of COVID-19 infection in older patients namely between 50 and 80 years old. The case fatality rate (CFR) was very high in critical cases that received MV namely 22/28 cases (78.57 %) (Table 4).

3.2 Risk Factors Associated with MV

The mortality rate namely case fatality rate (CFR %) was high overall, with 22/28 cases (78.57%) with critical COVID-19 dying. Of these, the majority (53.57 %, $n=15$) were afflicted with underlying co-morbidities, including 3 cases (10.71 %) each of diabetes mellitus or cardiovascular disorders as single co-morbidities, and 4 cases (14.28 %) with both diabetes mellitus and cardiovascular disorders together. 3 cases (10.71 %) had bronchial asthma, including 1 case (3.57%) with cardiovascular disorders as an additional co-morbidity, and 1 case (3.57 %)

had acute renal failure. Of 1 patients (3.57%) with underlying infections, the patient with HCV succumbed to COVID-19. On the other hand, 7 cases (28%) died without having any chronic diseases or co-infections (Fig. 1).

4. DISCUSSION

“Severe respiratory failure from COVID-19 pneumonia not responding to non-invasive respiratory support requires MV. Although ventilation can be a life-saving therapy, it can cause further lung injury if airway pressure and flow and their timing are not tailored to the respiratory system mechanics of the individual patient. The pathophysiology of COVID-19 infection can lead to a pattern of lung injury in patients with severe COVID-19 pneumonia typically associated with two distinct phenotypes, along a temporal and pathophysiological continuum, characterized by different levels of elastance, ventilation-to-perfusion ratio, right-to-left shunt, lung weight and recruitability. Understanding the underlying pathophysiology, duration of symptoms, radiological characteristics and lung mechanics at the individual patient level is crucial for the appropriate choice of MV settings to optimize gas exchange and prevent further lung injury. Previous study propose fundamental physiological and mechanical criteria for the selection of ventilation settings for COVID-19 patients in ICU. In particular, the choice of tidal volume should be based on obtaining a driving pressure < 14 cmH₂O, ensuring the avoidance of hypoventilation in patients with preserved compliance and of excessive strain in patients with smaller lung volumes and lower lung compliance. The level of PEEP should be informed by the measurement of the potential for lung recruitability, where patients with greater recruitability potential may benefit from higher PEEP levels. Prone positioning is often beneficial and should be considered early. The rationale for the proposed mechanical ventilation settings criteria is presented and discussed” [26].

“Early after establishment of MV, COVID-19 patients follow ARDS physiology, with compliance reduction related to the degree of hypoxemia, and inter-individually variable respiratory mechanics and recruitability” [27]. “COVID-19-ARDS is a subset of ARDS characterized overall by higher compliance and lung gas volume for a given PaO₂/FiO₂, at least when considered within the timeframe of our study” [28].

Table 1. National guideline of COVID-19 Critical Illness, Hodeidah, Yemen

Medicines Used	Critical
Paracetamol 1000 mg IV	three times a day (10-14 days)
Azithromycin 500 mg Tablet	once a day (6 days)
Ondansetron 8 mg IV	If necessary
Enoxaparin 60 mg Subq.	once a day (14 days)
Pantoprazole 40 mg IV	once a day (14 days)
Acetylcysteine IV 300 mg	three times a day (14 days)
Oseltamivir 75 mg Tablet	once dose (14 days)
Dexamethasone 10 – 20 mg IV	twice a day (10 days)
Piperacillin and Tazobactam 4.5 g IV	three times a day (10-14 days)
Moxifloxacin 400 mg IV	once a day (10 -14 days)
Meropenam 1 g IV	three times a day (10-14 days)
Vitamin D 50000 IU Tablet	once weekly (two weeks)
Vitamin C 1g IV	once a day (14 days)
Zink 300 mg Tablet	once dose (14 days)

Note:

- 1) Dose of dexamethasone was prescribed 20 mg in first five days and reduced into 10 mg in other five (first wave of COVID-19)
- 2) The doses of antibiotics were adjusted according to creatinine clearance. Antibiotic used to prevent the secondary infection and most cases had leukocytosis.
- 3) *Critical cases (acute respiratory distress syndrome "ARDS: respiratory failure, shock, or multi-organ system dysfunction).*

Table 2. Mechanical Ventilator (if needing) adjustment the parameters according the case

Variable	Setting
Ventilator mode	Volume assist – control
Tidal volume (ml kg ⁻¹)	6 (adjusted according to plateau pressure)
Plateau pressure (cm H ₂ O)	< 30
Rate (bpm)	6-35
I:E ratio	1:1 – 1:3
Oxygenation target	7.3 – 10.7
- PaO ₂	88-95
- SpO ₂	Set according to predetermined combinations
PEEP and FIO ₂	(PEEP range 5-24 cm H ₂ O)

[20,21]




Fig. 1. Risk factors of COVID-19 in first wave 2020 for patients undergo MV

Table 3. Mechanical ventilator variables for recovered case

Mechanical Ventilator Parameters	COVID-19 Patients
• RR	22
• Vt	380
• M-V	8.3 L
• FiO2	100
• PEEP	8
• I:E	1.2
• Flow rate,	50
• PIP	30
• Ti	1.3
• Ventilator mode	Simv_V

The MV variables included Respiratory rate (RR); Tidal volume (Vt); Fraction of inspired oxygen (FiO2); Minute volume (M-V); Time inspiration (Ti); Positive end-expiratory pressure (PEEP); inspiratory to expiratory ratio (I:E), Flow rate , Peak inspiratory pressure (PIP) , Plateau pressure (Pplat) , SaO₂, PaO₂, PaCo₂, pH, and ventilator modes that were adjusted .

Medications with MV:

- Midazolam 0.01-0.1 mg/kg
- Ketamine 2.5 – 15 mcg/kg/min
- Propofol 25-100 mcg/kg/min
- Atracurium 0.005-0.01 mg/kg/min
- Fentanyl 1-2 mcg/kg/h
- Morphine 0.05-0.2 mg/kg
- Pethadine 0.25 mg [22-25]

Table 4. Sex – age and outcome of critical illness with COVID-19 infection undergo MV

Sex/Age	Mortality			Recovery			Total (n)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<15	1	0	1	0	1	1	1	1	2
15-29	2	0	2	1	0	1	3	0	3
30 -49	0	0	0	1	0	1	1	0	1
50-59	6	2	8	2	1	3	8	3	11
60+	8	3	11	0	0	0	8	3	11
	17	5	22	4	2	6	21	7	28
Total	22			6			28		

“The most commonly used analgesics and sedatives for all patients admitted to the ICU were pethidine (26.14%) and midazolam (32.18%), respectively. Sedatives and analgesics were more commonly used in mechanically ventilated patients. Among analgesics, the usage rate of pethidine and morphine decreased, whereas the usage rate of fentanyl and remifentanyl increased. Among sedatives, the usage rate of benzodiazepine decreased, whereas the usage rate of propofol increased [24]. There was discordance between current usage of analgesics and sedatives and the recommended usage stipulated by ICU guidelines. However, the trend of drug usage is changing to match the guidelines, which recommend maintenance of light sedation using an analgesia-based regimen and usage of short-

acting drugs for routine monitoring of pain, agitation, and delirium in ICU care” [25].

“For a similar initial oxygenation, COVID-19 ARDS initially differs from classical ARDS by a higher CRS, dissociated from oxygenation. CRS become similar for patients remaining on mechanical ventilation during the first week of evolution, but oxygenation becomes lower in COVID-19 patients” [29]. “The compliance of the respiratory system is similar between COVID ARDS and non-COVID ARDS when calculated at the same PEEP level and while taking into account patients’ anthropometric characteristics” [30].

“Old age and co-morbidity with chronic diseases namely diabetes, hypertension, cardiac disorder,

and asthma that may be contributing factors to excess deaths among COVID-19 patients [9-10]. On the other hand, the co-infections with other infections like vector – borne diseases dengue, malaria, and west – Nile virus is of high concern in Hodeidah, Yemen” [31-35].

On the other hand, study in Yemen reported that 9.7 % of COVID-19 patients needed ICU admission (severe and critical illness) [10] that is similar with previous study that reported range of ICU admission with COVID-19 infection from 9.4 to 45.9% [36]. In addition, the finding reported that the CFR (%) of admitted patients (critical illness) to ICU was 23/49 cases (46.9%). On the other mean, 28/49 cases (57.14 %) in isolation department who had received MV and 22/28 cases (78.57 %) of those died and this result agreed with other countries. “In China, Wuhan, mortality rates among those admitted to ICUs ranged from 52 to 62% and increased to 86–97% among those requiring invasive MV. In United Kingdom, 67 % of those who had received MV died. In the United States, indicated that 50–67% of patients admitted to the ICU and 71–75% of those receiving invasive MV died” [37].

5. CONCLUSION

The impact of MV on mortality outcome was very high in Hodeidah, Yemen. Also, the COVID-19 has complications that increases the mortality rate. In addition, the old age, chronic diseases and co-infection may be contributing factors to excess mortality among COVID-19 patients. On the other hand, a greater emphasis toward training in best practice of MV and ICU medicine namely respiration therapy, are needed to help overcome these challenges.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

CONSENT

As per international standards or university standards, Participants' written consent has been collected and preserved by the authors. The raw data are secured in the Center of Tropical Medicine and Infectious Diseases (CTMID), Al-Thawara Public Hospital Authority, Hodeidah, Yemen.

ETHICAL APPROVAL

The studies involving human participants were reviewed and approved by the Ethics Committee of the Center for Tropical Medicine and Epidemiology Studies, Hodeidah University (CTMES – HU), Hodeidah, Yemen.

ACKNOWLEDGEMENTS

The authors would like to thank the supervisor and medical staff of the COVID-19 isolation department, Molecular Biology Unit, Center of Tropical Medicine and Infectious Diseases (CTMID), AL Thawara Public Hospital Authority, Hodeidah, Yemen for their fruitful assistance.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Butler MJ, Best JH, Mohan SV, Jonas JA, Arader L, Yeh J. Mechanical ventilation for COVID-19: Outcomes following discharge from inpatient treatment. *PloS one*, (2023). 18(1), e0277498. Available: <https://doi.org/10.1371/journal.pone.0277498>
2. Nolley EP, Sahetya SK, Hochberg CH, Hossen S, Hager DN, Brower RG, Stuart EA, Checkley W. Outcomes Among Mechanically Ventilated Patients with Severe Pneumonia and Acute Hypoxemic Respiratory Failure From SARS-CoV-2 and Other Etiologies. *JAMA network open*. 2023;6(1):e2250401. Available: <https://doi.org/10.1001/jamanetworkopen.2022.50401>
3. Roedl K, Jarczak D, Thasler L, Bachmann M, Schulte F, Bein B, Weber CF, Schäfer U, Veit C, Hauber HP, Kopp S, Sydow K, de Weerth A, Bota M, Schreiber R, Detsch O, Rogmann JP, Frings D, Sensen B, Burdelski C, Kluge S. Mechanical ventilation and mortality among 223 critically ill patients with coronavirus disease 2019: A multicentric study in Germany. *Australian critical care: official journal of the Confederation of Australian Critical Care Nurses*. 2021;34(2):167–175. Available: <https://doi.org/10.1016/j.aucc.2020.10.009>
4. Krynytska I, Marushchak M, Birchenko I, Dovgalyuk A, Tokarsky O. COVID-19-

- associated acute respiratory distress syndrome versus classical acute respiratory distress syndrome (a narrative review). *Iranian journal of microbiology*. 2021;13(6):737–747. Available:<https://doi.org/10.18502/ijm.v13i6.8072>
5. López-Hernández D. Clinical Presentation and Factors Associated to COVID-19 Disease in Mexican Patients. *Curr. J. Appl. Sci. Technol*. 2022;41(1):40-58. [accession: 2024 Jun. 5]; Available:<https://journalcjust.com/index.php/CJAST/article/view/3804>
 6. Pai MYB, Park ASH, Chen BFL, Toma JT, Pai HJ. Health Complications in Patients Recovering from COVID-19: A Narrative Review of Post-COVID Syndrome. *J. Adv. Med. Med. Res*. 2021;33(10):115-29. [accession: 2024 Jun. 5]; Available:<https://www.journaljammr.com/index.php/JAMMR/article/view/4011>
 7. Hua J, Qian C, Luo Z, Li Q, Wang F. Invasive mechanical ventilation in COVID-19 patient management: the experience with 469 patients in Wuhan. *Critical Care*. 2020;24:1-3.
 8. Cronin JN, Camporota L, Formenti F. Mechanical ventilation in COVID-19: A physiological perspective. *Experimental physiology*. 2022;107(7):683-93.
 9. Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, Wu Y, Zhang L, Yu Z, Fang M, Yu T, Wang Y, Pan S, Zou X, Yuan S, Shang Y. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *The Lancet. Respiratory medicine*. 2020;8(5):475–481. Available:[https://doi.org/10.1016/S2213-2600\(20\)30079-5](https://doi.org/10.1016/S2213-2600(20)30079-5)
 10. AL Kamarany MA, Al-Masrafi I, Ghouth AB, Suhail K, Majam A, Zuhairy A, Alabsi E. Outbreak Investigation: First Ten COVID – 19 Infection Related Deaths in Hodeidah, Yemen. *Asian Journal of Research in Infectious Diseases*. 2023;13(3):19–27. Available:<https://doi.org/10.9734/ajrid/2023/v13i3268>
 11. AL-Kamarany MA, Suhail KA, Majam AS, Abdulabari Alabsi E, Hamoud Dowbalah M, Mohammed Zohairy, A. Epidemiological and Clinical Features of COVID-19 in Hodeidah, Yemen. *International Journal of TROPICAL DISEASE & Health*. 2021;42(21):28–40. Available:<https://doi.org/10.9734/ijtdh/2021/v42i2130550>
 12. AL Kamarany MA, Abdulkarim T, & Nasser M. Biomolecular Features of COVID-19 in Hodeidah, Yemen. *Asian Journal of Biochemistry, Genetics and Molecular Biology*. 2023;14(3):29–38. Available:<https://doi.org/10.9734/ajbgmb/2023/v14i3317>
 13. Al-Kamarany MA, Abdulkarim T. Immunological Features of COVID-19 in Hodeidah, Yemen. *Asian Journal of Immunology*. 2023;6(1):76–84. Available:<https://journalaji.com/index.php/AJI/article/view/97>
 14. World Health Organization (WHO). Operation consideration for case management of COVID-19 Health Facility;2020.
 15. Ministry of Public Health and Population (MOPHP) of, Syrian National Guideline for Control of Coronavirus Disease. 2019 (COVID-19);2020.
 16. Suhail FA, Al Kamarany MA. Radiological Features of COVID-19 Patients in Hodeidah, Yemen. *Asian Journal of Research in Infectious Diseases*. 2021;8(4):117–127. Available:<https://doi.org/10.9734/ajrid/2021/v8i430256>
 17. Balkam FA, Al-Kamarany MA. Hematological and coagulation features of COVID-19 patients in Hodeidah, Yemen. *International Journal of Pathogen Research*. 2023;12(6):46–54. Available:<https://doi.org/10.9734/ijpr/2023/v12i6252>
 18. AL-Kamarany MA, Balkam FA. Biochemical Features of COVID-19 Patients in Hodeidah, Yemen. *Asian Journal of Research in Biosciences*. 2024; 6(1):96–103. Available:<https://globalpresshub.com/index.php/AJORIB/article/view/1997>
 19. AL-Kamarany MA, Alkadi HO, Almadani MY, Alsharma ZA, Al-Salehy RM, Albaser NA. Pharmacological and Therapeutic Features of COVID - 19 Infection in Hodeidah, Yemen. *Journal of Advances in Medical and Pharmaceutical Sciences*. 2024;26(5):18–38. Available:<https://doi.org/10.9734/jamps/2024/v26i5683>
 20. Fontela PC, Prestes RB, Forgiarini LA, Jr, Friedman G. Variable mechanical

- ventilation. *Ventilação mecânica variável. Revista Brasileira de terapia intensiva.* 2017;29(1):77–86.
Available:<https://doi.org/10.5935/0103-507X.20170012>
21. Sean M. Hickey; Al O. Giwa. Mechanical Ventilation, Last Update; 2023.
Available:<https://www.ncbi.nlm.nih.gov/books/NBK539742/>
 22. Pearson SD, Patel BK. Evolving targets for sedation during mechanical ventilation. *Current opinion in critical care.* 2020; 26(1):47–52.
Available:<https://doi.org/10.1097/MCC.0000000000000687>
 23. Umunna BP, Tekwani K, Barounis D, Kettaneh N, Kulstad E. Ketamine for continuous sedation of mechanically ventilated patients. *Journal of emergencies, trauma, and shock.* 2015; 8(1):11–15.
Available:<https://doi.org/10.4103/0974-2700.145414>
 24. Kim HH, Choi SC, Ahn JH, Chae MK, Heo J, Min YG. Analysis of trends in usage of analgesics and sedatives in intensive care units of South Korea: A retrospective nationwide population-based study. *Medicine.* 2018;97(35):e12126.
Available:<https://doi.org/10.1097/MD.00000000000012126>
 25. Aoki Y, Kato H, Fujimura N, Suzuki Y, Sakuraya M, Doi M. Effects of fentanyl administration in mechanically ventilated patients in the intensive care unit: a systematic review and meta-analysis. *BMC anesthesiology.* 2022;22(1):323.
Available:<https://doi.org/10.1186/s12871-022-01871-7>
 26. Cronin JN, Camporota L, Formenti F. Mechanical ventilation in COVID-19: A physiological perspective. *Experimental physiology.* 2022;107(7):683–693.
Available:<https://doi.org/10.1113/EP089400>
 27. Grieco DL, Bongiovanni F, Chen L, Menga LS, Cutuli SL, Pintaudi G, Carelli S, Michi T, Torrini F, Lombardi G, Anzellotti GM, De Pascale G, Urbani A, Bocci MG, Tanzarella ES, Bello G, Dell'Anna AM, Maggiore SM, Brochard L, Antonelli M. Respiratory physiology of COVID-19-induced respiratory failure compared to ARDS of other etiologies. *Critical care (London, England).* 2020;24(1):529.
Available:<https://doi.org/10.1186/s13054-020-03253-2>
 28. Chiumello D, Busana M, Coppola S, Romitti F, Formenti P, Bonifazi M, Pozzi T, Palumbo MM, Cressoni M, Herrmann P, Meissner K, Quintel M, Camporota L, Marini JJ, Gattinoni L. Physiological and quantitative CT-scan characterization of COVID-19 and typical ARDS: a matched cohort study. *Intensive care medicine.* 2020;46(12):2187–2196.
Available:<https://doi.org/10.1007/s00134-020-06281-2>
 29. Beloncle F, Studer A, Seegers V, Richard JC, Desprez C, Fage N, Merdji H, Pavlovsky B, Helms J, Cunat S, Mortaza S, Demiselle J, Brochard L, Mercat A, Meziani F. Longitudinal changes in compliance, oxygenation and ventilatory ratio in COVID-19 versus non-COVID-19 pulmonary acute respiratory distress syndrome. *Critical care (London, England).* 2021;25(1): 248.
Available:<https://doi.org/10.1186/s13054-021-03665-8>
 30. Fusina F, Albani F, Crisci S, Morandi A, Tansini F, Beschi R, Rosano A, Natalini G. Respiratory system compliance at the same PEEP level is similar in COVID and non-COVID ARDS. *Respiratory research.* 2022;23(1):7.
Available:<https://doi.org/10.1186/s12931-022-01930-0>
 31. Al Kamarany MA, Majam A, Suhail K, Zuhairy A, Alabsi E. Coronavirus Disease 2019 – Dengue Fever Coinfection: A case report. *International Journal of Pathogen Research.* 2023;12(4):27–32.
Available:<https://doi.org/10.9734/ijpr/2023/v12i4234>
 32. Alahdal M, Al-Shabi J, Ogaili M, Abdullah QY, Alghalibi S, Jumaan AO, Al-Kamarany MA. Detection of dengue fever virus serotype – 4 by using One-Step Real-Time RT-PCR in Hodeidah, Yemen. *Microbiology Research Journal International.* 2016;14(6):1–7.
DOI:<https://doi.org/10.9734/BMRJ/2016/24380>
 33. Al-Areeqi A, Alghalibi S, Yusuf Q, Al-Masrafi I, Al-Kamarany MA. Epidemiological characteristic of malaria coinfecting with dengue fever in Hodeidah, Yemen. *International Journal of TROPICAL DISEASE & Health,* 2020;40(3):1–10.

- DOI:<https://doi.org/10.9734/ijtdh/2019/v40i330230>
34. Yusuf QA, Ogaili M, Alahdal M, Amood Al Kamarany M. Dengue fever infection in Hodeidah, Yemen risk factors and socioeconomic indicators. *British Biomedical Bulletin*. 2015;3(1):58–65.
35. Yusuf Q, Al-Masrafi I, Al-Mahbashi A, Al-Areeqi A, Al-Kamarany MA, & Khan AS. First evidence of West Nile Virus in Hodeidah, Yemen: Clinical and epidemiological characteristics. *International Journal of TROPICAL DISEASE & Health*. 2019;38(4):1-9.
- DOI:<https://doi.org/10.9734/ijtdh/2019/v38430190>
Available:<https://journalijtdh.com/index.php/IJTDH/article/view/30190>
36. Paolo I, Nicola M, Elio A, Guido R, Mario B and Donata G. COVID-19 mortality and ICU admission: the Italian experience. *Critical Care* 2020;24:228.
37. Auld Sara C et al. ICU and Ventilator Mortality Among Critically Ill Adults with Coronavirus Disease 2019. *Critical Care Medicine*. 2020;48:9:e799-e804.
DOI: 10.1097/CCM.0000000000004457
DOI: 10.1148/radiol.2020201473

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://prh.globalpresshub.com/review-history/1612>