



Effect of Phosphorus, Sulphur and Gibberellic Acid on Growth, Root Nodules and Yield of Soybean [*Glycine max* (L.) Merrill]

Bahnu Pratap Ghasil ^a, Harphool Meena ^{b*}, Manoj ^c,
Rajendra K. Yadav ^b, Monika Choudhary ^d,
Govind Ram Yadav ^c, Dhuni Lal Yadav ^b,
Shankar Lal Yadav ^b and R. K. Meena ^a

^a College of Agriculture, Agriculture University, Kota, Rajasthan, India.

^b Agricultural Research Station, Agriculture University, Kota, Rajasthan, India.

^c Department of Soil Science and Agricultural Chemistry, SKNAU, Jobner, Jaipur, Rajasthan, India.

^d Department of Agronomy, Rajasthan College of Agriculture, Udaipur, Rajasthan, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i242735

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96493>

Original Research Article

Received: 23/10/2022

Accepted: 30/12/2022

Published: 31/12/2022

ABSTRACT

An experiment was conducted during *kharif* season 2020 at Instructional Farm, College of Agriculture, Umedganj, Kota. The experiment comprised twenty-four treatment combinations, having four levels of phosphorus *viz.*, 0, 20, 40 and 60 kg ha⁻¹, three levels of sulphur *viz.*, 15, 30 and 45 kg ha⁻¹ and foliar spray of gibberellic acid *viz.*, no spray and spray of GA₃ @ 75 ppm laid out

*Corresponding author: E-mail: hpagron@rediffmail.com;

in sub-sub split-plot design with three replications. Results showed that application of 60 kg P₂O₅ ha⁻¹ had significant effect on plant height, dry matter accumulation plant⁻¹, branches plant⁻¹ at 30, 60 DAS and at harvest, root nodules plant⁻¹ and their dry weight at 45 DAS while root nodules plant⁻¹ and their dry weight at 45 DAS and seed yield (1956 kg ha⁻¹) over application of 40, 20 kg P₂O₅ ha⁻¹ and control. Significantly higher seed yield (1956 kg ha⁻¹) was recorded under application of 60 kg P₂O₅ ha⁻¹ over application of 20 kg P₂O₅ ha⁻¹ and control. However, it was found at par with application of 40 kg P₂O₅ ha⁻¹. Application of 45 kg sulphur ha⁻¹ had significantly higher plant height (cm) at 60 DAS and at harvest, dry matter accumulation (g plant⁻¹) at 30, 60 and at harvest stages, branches plant⁻¹ at 30, 60 and at harvest stage, number of root nodules and their dry weight (g) and seed yield (1742 kg ha⁻¹) which was found at par with application of 30 kg sulphur ha⁻¹ over 15 kg sulphur ha⁻¹. Application of gibberellic acid @ 75 ppm as foliar spray gave significantly higher plant height (cm) at 60 DAS and at harvest stage, dry matter accumulation (g plant⁻¹) at 60 DAS and at harvest, branches plant⁻¹ at 30, 60 DAS and harvest and seed yield (1770 kg/ha) over control.

Keywords: Growth; root nodules; seed yield; phosphorus; sulphur; gibberellic acid.

1. INTRODUCTION

In India oilseed crops constitute the second largest agricultural produce, next to food grain and these crops are the important sources of fats and oils. "The oil and economic end product of oilseed crop is an integral part of human diet. Beside the dietary needs, the vegetable edible oil has numerous mechanical, industrial, medicinal and therapeutic uses too. Soybean has paramount importance in human and animal nutrition because it is a major source of edible vegetable oil and high protein feed as well as food in the world. Soybean is considered as miracle crop because it contains 38 - 42 per cent good quality protein, 23 per cent carbohydrates, 18-20 per cent oil, rich in poly unsaturated fatty acids, a good amount of minerals and vitamins especially B-complex and tocopherols. It provides high amounts of phyto chemicals and good quality dietary fibre which enables to protect human body against cancers and diabetes" [1].

"India ranks fifth in the world in area and production after USA, Brazil, Argentina and China. Soybean has emerged as an important oilseed crop in India. On a national basis, soybean occupied an area of 12.09 million ha with production and productivity of 11.22 metric tonnes and 928 kg ha⁻¹, respectively" [2]. "Soybean is grown as a major oilseed crop mainly in south-eastern parts of Rajasthan during *kharif* season. It covers 1.12 million ha with annual production and productivity of 0.52 metric tonnes and 469 kg ha⁻¹ respectively in the state [2]. Six districts namely Kota, Bundi, Baran, Jhalawar, Sawai madhopur and Karauli of state come under the jurisdiction area of Agriculture University, Kota. Soybean crop has an area of 7.26 lakh ha with annual production 3.43 lakh

tones and average productivity is 636 kg ha⁻¹ [3], which is quite less than its potential yield owing to various stresses during the growing season".

"Plants require phosphorus for growth throughout their life cycle, especially during the early stages of growth and development. In soybean, the demand for phosphorus is the greatest during pod and seed development stage where more than 60 per cent of phosphorus end up in the pods and seeds" [4]. Its uptake and utilization by soybean are essential for ensuring proper nodule formation and improving yield and quality of the crop. Sulphur plays a pivotal role in various plant growth and development processes being a constituent of sulphur containing amino acids, cystine and methionine, and other metabolites *viz.*, lutein and phytochelators.

"Sulphur is used as soil amendment for amelioration, as plant nutrient for increasing yield and quality of crop produce, as chemical agent to acidulate other nutrient and pesticides" [5]. "A higher susceptibility of crops to certain diseases was observed in sulphur deficient soils" [6]. It is also implicated in oil biosynthesis in soybean seed [7]. "Still the studies on effect of sulphur in soybean are very meagre. The sulphur is required in high amount by the oilseeds and hence has been identified as key nutrient responsible for high production. Plant growth regulators are known to enhance the source-sink relationship and stimulate the translocation of photo-assimilates there by helping in improve the physiological efficiency, photosynthetic ability, effective partitioning of assimilates from source to sink and ultimately enhance productivity of the soybean crop" [8]. "Gibberellic acid constitutes a group of tetracyclic diterpenoids, involved in plant growth and development. A well-known phyto-hormone has numerous physiological effects on

plants including seed germination, growth, stem elongation, leaf expansion, photosynthesis, flowering, cell expansion and also increase in activities of many key enzymes like carbonic anhydrase, nitrate reductase in field crops" [9].

2. MATERIALS AND METHODS

An experiment was conducted during *kharif* season 2020 at Instructional Farm, College of Agriculture, Umedganj, Kota (Rajasthan), which is situated in South-Eastern part of Rajasthan. In Rajasthan, this region falls under Agro-climatic zone V B (Humid South eastern Plains) of Rajasthan. This zone possesses typical sub-tropical conditions with maximum and minimum temperatures ranged between 34.2°C to 38.0°C and 18.6°C to 24.0 °C during *Kharif*, 2020. The total amount of rainfall received during crop growing was 551 mm. The soil of the experimental site was clay loam in texture, slightly saline in reaction. The experimental soil was medium in available nitrogen (264 kg ha⁻¹) and phosphorus (21.7 kg ha⁻¹) while high in potassium (388 kg ha⁻¹) and sufficient in DTPA extractable micronutrients with pH (7.61) and EC (0.52 dS m⁻¹). The source of nutrients applied were urea for nitrogen, DAP for phosphorus and mutate of potash for potassium.

The experiment consisted of twenty-four treatment combinations including four levels of phosphorus (control, 20, 40 and 60 kg/ha) allocated in main plots, three levels of sulphur (15, 30 and 45 kg/ha) in sub plots and two levels of foliar application of gibberellic acid (foliar spray of GA3 @ 75 ppm and no spray of GA3) in sub-sub plots were under taken in sub-sub split plot design with replicated thrice. Data on growth parameters like plant height, dry matter accumulation, branches plant⁻¹, root nodules, nodules dry weight and seed yield were recorded as per standard procedures. For recording of pre and post-harvest observations, five plants were randomly selected for each plot and tagged with labels for various observations. Plant samples were collected from each of the plots. "The data were statistically analysed by adopting an appropriate method of standard analysis of variance" [10].

3. RESULTS AND DISCUSSION

3.1 Effect of Phosphorus on Growth Parameters

Data presented in Table 1 revealed that plant population of soybean at 30 DAS and at harvest

was not significantly affected by any of the treatment combination. Thus, plant population was almost uniform in all the treated plots. The phosphorus fertilization failed to bring perceptible variation in plant height at 30 DAS, while variation in plant height due to difference level of fertilizer was significant at 60 DAS and at harvest. Significantly tallest plant height (55.97 cm) at 60 DAS and (75.93 cm) at harvest was recorded with the application of 60 kg P₂O₅ ha⁻¹ which remained statistically at par with 40 kg P₂O₅ ha⁻¹ and over 20 kg P₂O₅ ha⁻¹ and control. Application of 60 kg P₂O₅ ha⁻¹ gave significantly higher dry matter accumulation (3.56, 12.02 and 16.63 g plant⁻¹) at 30, 60 DAS and at harvest, respectively which was closely followed by application of 40 kg P₂O₅ ha⁻¹ (3.15, 10.68 and 15.20 g plant⁻¹) over application of 20 kg P₂O₅ ha⁻¹ and control. The higher number of branches (2.98, 4.49 and 5.15 plant⁻¹) at 30, 60 DAS and at harvest was recorded with the application of phosphorus 60 kg ha⁻¹ over application of 20 kg phosphorus ha⁻¹ and control. However, it was found at par with application of phosphorus 40 kg ha⁻¹ (2.69, 4.01 and 4.67 plant⁻¹) at 30, 60 DAS and at harvest. The maximum number of root nodules plant⁻¹ (52.02) at 45 DAS was recorded with application of 60 kg P₂O₅ ha⁻¹ which was closely followed by 40 kg P₂O₅ ha⁻¹ over control. Significantly higher dry weight of root nodules (84.44 mg plant⁻¹) at 45 DAS was recorded with the application of 60 kg P₂O₅ ha⁻¹ which was closely followed by application of 40 kg P₂O₅ ha⁻¹ dry weight of root nodules (80.02 mg plant⁻¹) over application of 20 kg P₂O₅ ha⁻¹ and control (Table 2).

"Plant height may be increased due to uptake of nitrogen and phosphorus by the plants, which was made available through nitrogen fixation and phosphorus solubilisation by the beneficial microorganisms" [11]. "Phosphorus promotes root growth, cell formation, leaf development, seed formation and accelerates early maturity of crop which may result to increment in branches, root nodules of plant" [12]. "This might be due to the fact that application of phosphorus results profuse growth of roots which ultimately resulted formation of a greater number of large size root nodules" [13]. "Each increment in phosphorus fertilizer levels recorded significant variations in leaf area index and green leaves plant⁻¹ at 45 DAS. This reveals that increasing phosphorus level enhanced the soil phosphorus availability and consequently it's uptake by soybean crop plants which led to higher size of photosynthetic apparatus. This results is supported by significantly higher chlorophyll content and leaf

area index at different growth stages of the crop by many workers including" [14].

3.2 Effect of Phosphorus on Yield

A reference data was recorded during experimentation and data presented in Table 2. Data further showed that various level of phosphorus fertilizer significantly influenced seed yield of soybean. The maximum seed yield (1956 kg ha⁻¹) was recorded with the application of 60 kg P₂O₅ ha⁻¹ over application of 20 kg P₂O₅ ha⁻¹ and control seed yield (1559 and 1176 kg ha⁻¹). However, it was found at par with application of 40 kg P₂O₅ ha⁻¹ seed yield (1877 kg ha⁻¹). "Phosphorous application accelerated the production of photosynthates and their translocation from source to sink, which ultimately gave the higher values of yield contributing characters. Increase in yield contributing characters" has also been reported by Meena et al. [15] and Kumar et al. [16]. "This was mainly due to fact that the better availability of nitrogen and phosphorus caused well developed root system having higher nitrogen fixing capacity resulting better growth and development of plants and better translocation of photosynthates towards sink, even use of single or combination of fertilizers might be much advantageous for farmers" [17].

3.3 Effect of Sulphur on Growth Parameters

Data presented in Table 1 revealed that plant population of soybean at 30 DAS and at harvest was not significantly affected by any of the treatment combination. Thus, plant population was almost uniform in all the treated plots. Data further indicated that plant height at 30 DAS was remained significantly unaffected with application of sulphur. Significantly highest plant height (53.05 cm) at 60 DAS and (73.57 cm) at harvest was recorded with the application of 45 kg sulphur ha⁻¹ which was closely followed by 30 kg sulphur ha⁻¹ at 60 DAS (49.88 cm) and at harvest (70.45 cm). Significantly highest dry matter accumulation (3.15, 10.58 and 15.16 g plant⁻¹) at 30, 60 DAS and at harvest was recorded with the application of 45 kg sulphur ha⁻¹, which was closely followed by application of 30 kg sulphur ha⁻¹ (2.90, 9.99 and 13.87 g plant⁻¹) at 30, 60 DAS and at harvest over application of 15 kg sulphur ha⁻¹. Significantly higher number of branches (2.78, 4.0 and 4.83 plant⁻¹) at 30, 60 DAS and at harvest was recorded with the application of 45 kg sulphur ha⁻¹ which was closely followed by application of 30 kg sulphur

ha⁻¹ (2.54, 3.75 and 4.44 branches plant⁻¹). Minimum number of branches plant⁻¹ was recorded under application of 15 kg sulphur ha⁻¹ Table 2. Application of 45 kg sulphur ha⁻¹ was recorded significantly higher root nodules plant⁻¹ (49.09) at 45 DAS over application of sulphur 15 kg ha⁻¹ (42.49) root nodules plant⁻¹. However, it was found at par with the application of 30 kg sulphur ha⁻¹ root nodules plant⁻¹ (46.30). Application of 45 kg sulphur ha⁻¹ was recorded significantly higher dry weight of root nodules (80.84 mg plant⁻¹) at 45 DAS over application of 15 kg sulphur ha⁻¹. However, it was found on par with the application of 30 kg sulphur ha⁻¹ dry weight of root nodules (75.33 mg plant⁻¹).

"This may be due to better root development and profuse nodulation on account of increased *rhizobial* activity in the rhizosphere under sulphur and bio fertilizers availability. This finally resulted in the formation of bolder and more number of root nodules. The positive response of sulphur on nodulation" was also observed by Watimongla et al. [18]. "The plant height and branches improved by sulphur alone or combined with nitrogen whereas, nitrogen alone decreased number of pods plant⁻¹ thus showing non-significant (P≤0.05) variation in grain yield as compared to control. The results corroborate the findings with sulphur application of 40 kg ha⁻¹ enhanced the plant height and branches in soybean" [19]. The sulphur fertilizer levels recorded significant variations in leaf area index and green leaves plant⁻¹ at 45 DAS. This reveals that increasing sulphur level enhanced availability of sulphur in soil and consequently it's mining by soybean crop plants which led to higher size of photosynthetic apparatus. This statement is endorsed by significantly higher chlorophyll content and leaf area index at different growth stages of the crop. The increase in green leaves plant⁻¹ and leaf area index with sulphur levels has been ascribed to more dry matter accumulation, this might be due to high accumulation of net photosynthates. The results obtained are consistent with findings reported by Meena et al. [20].

3.4 Effect of Sulphur on Yield

Data was recorded during experimentation and data presented in Table 2 showed that various level of phosphorus fertilizer significantly influenced seed yield of soybean. The maximum seed yield (1742 kg ha⁻¹) was recorded with the application of 45 kg sulphur ha⁻¹ over application of 15 kg sulphur ha⁻¹ seed yield (1500 and 1176 kg ha⁻¹). However, it was found at par with

Table 1. Effect of phosphorus, sulphur and gibberellic acid on growth parameters of soybean

Treatments	Plant population (mrl ⁻¹)		Plant height (cm)			Plant dry matter accumulation (g plant ⁻¹)		
	At 30 DAS	At harvest	At 30 DAS	At 60 DAS	At harvest	At 30 DAS	At 60DAS	At harvest
A. Phosphorus (kg ha⁻¹)								
0	10.31	10.20	21.2	43.40	64.15	2.12	7.21	10.13
20	10.35	10.25	21.96	48.08	67.32	2.74	9.08	13.14
40	10.42	10.31	22.62	52.83	72.45	3.15	10.68	15.20
60	10.49	10.40	23.18	55.97	75.93	3.56	12.02	16.63
SEm±	0.24	0.23	0.50	1.02	1.15	0.06	0.17	0.30
CD at 5%	NS	NS	NS	3.54	3.98	0.20	0.61	1.05
B. Sulphur (kg ha⁻¹)								
15	10.33	10.22	21.54	47.27	65.82	2.62	8.67	12.61
30	10.40	10.30	22.42	49.88	70.45	2.90	9.99	13.87
45	10.44	10.36	22.76	53.05	73.57	3.15	10.58	15.16
SEm±	0.12	0.14	0.38	1.00	1.01	0.05	0.20	0.16
CD at 5%	NS	NS	NS	3.02	3.05	0.15	0.62	0.49
C. Gibberellic acid								
No spray	10.37	10.20	21.87	48.16	67.45	2.83	9.44	13.13
GA ₃ @ 75 ppm	10.41	10.38	22.61	51.96	72.45	3.95	10.05	14.42
SEm±	0.13	0.13	0.27	0.78	1.02	0.04	0.14	0.15
CD at 5%	NS	NS	NS	2.28	2.99	NS	0.42	0.44

Table 2. Effect of phosphorus, sulphur and gibberellic acid on growth and yield of soybean

Treatments	Branches (plant ⁻¹)			Nodules at 45 DAS (No. plant ⁻¹)	Nodules dry weight at 45 DAS (mg plant ⁻¹)	Seed yield (kg ha ⁻¹)
	At 30 DAS	At 60 DAS	At harvest			
A. Phosphorus (kg ha⁻¹)						
0	1.92	2.85	3.65	38.35	64.89	1176
20	2.37	3.45	4.18	44.21	73.36	1559
40	2.69	4.01	4.67	49.26	80.02	1877
60	2.98	4.49	5.15	52.02	84.44	1956
SEm±	0.05	0.05	0.11	0.69	1.23	37.28
CD at 5%	0.17	0.20	0.38	2.39	4.28	129.00
B. Sulphur (kg ha⁻¹)						
15	2.14	3.35	3.95	42.49	70.87	1500
30	2.54	3.75	4.45	46.30	75.33	1684
45	2.78	4.00	4.83	49.09	80.84	1742
SEm±	0.04	0.06	0.09	0.80	0.97	30.87
CD at 5%	0.14	0.19	0.28	2.40	2.91	92.57
C. Gibberellic acid						
No spray	2.41	3.53	4.28	45.14	74.53	1592
GA ₃ @ 75 ppm	2.56	3.87	4.54	46.78	76.83	1693
SEm±	0.04	0.06	0.05	0.68	0.97	25.77
CD at 5%	NS	0.19	0.17	NS	NS	75.23

application of 30 kg sulphur ha⁻¹ seed yield (1684 kg ha⁻¹). "The yield increased under sulphur fertilization might be ascribed to increased pods plant⁻¹ and seeds pod⁻¹ with heavier seeds. Thus, significant improvement in yield obtained under sulphur fertilization seems to have resulted owing to increased concentration of sulphur in various parts of plant that helped maintain the critical balance of other essential nutrients in the plant and resulted in enhanced metabolic processes. Vyas et al. [21] also noticed increased yield of soybean with application of sulphur. Sulphur plays a vital role in improving vegetative structure for nutrient absorption, strong sink strength through development of reproductive structures and production of assimilates to fill economically important sink" [22]. The seeds pod⁻¹ and seed yield improved by sulphur alone or combined with nitrogen whereas, nitrogen alone decreased number of pods plant⁻¹ thus showing non-significant variation in seed yield as compared to control. The results corroborate the findings with application of 40 kg sulphur ha⁻¹ enhanced the pod plant⁻¹ and test weight (g) in black gram [23].

3.5 Effect of Gibberellic acid Growth Parameters

Results revealed that in Table 1 the plant population of soybean was not significantly affected by foliar spray of GA₃. Application of foliar spray of GA₃ @ 75 ppm failed to bring about significant variation in plant height over control at 30 DAS, while the effect of application of foliar spray of GA₃ @ 75 ppm was significant (51.96 cm) at 60 DAS and (72.45 cm) at harvest over no spray. A critical examination of data shows that the foliar spray of GA₃ @ 75 ppm failed to bring about significant variation in dry matter accumulation over control at 30 DAS. The significantly highest dry matter accumulation (10.05 and 14.42 g plant⁻¹) at 60 DAS and at harvest was recorded with the application of foliar spray of GA₃ @ 75 ppm over control. The given data (Table 2) further indicated that application of foliar spray of GA₃ @ 75 ppm failed to bring about significant variation in number of branches plant⁻¹ over control at 30 DAS. Significantly highest branches at 60 DAS and at harvest was recorded with the application of foliar spray of GA₃ @ 75 ppm (3.87 and 4.54 plant⁻¹), respectively over control. Application of foliar spray of GA₃ @ 75 ppm and no spray of GA₃ was did not significantly affected root nodules plant⁻¹. The dry weight of root nodules was not significantly affected with the application of foliar spray of GA₃ @ 75 ppm and no spray.

Gibberellic acid is the most widely used plant growth regulator which increases stem elongation along with plant height, growth, dry matter accumulation as well as yield in various crops [24] and [25]. However, very few works have been done on the application of gibberellic acid on french bean in Bangladesh [26].

3.6 Effect of Gibberellic Acid on Yield

Data further (Table 2) indicated that application of foliar spray of GA₃ @ 75 ppm was gave significantly higher seed yield (1770 kg ha⁻¹) over no spray of gibberellic acid seed yield (1514 kg ha⁻¹) of soybean. Foliar application of plant growth regulators strengthened physiological relationship between source and sink resulting in effective partitioning and translocation of photosynthates from leaves to seeds within the plant. The application of growth regulators increased the test weight of soybean. This might be due to mark increased the vegetative growth, photosynthetic pigment which could increase in photosynthesis and increased the seed weight. The similar observation reported by Sapkal et al. [27] the application of GA₃ @ 50 ppm increased seed weight in soybean. The foliar application of GA₃ increased the test weight [28] and [29].

4. CONCLUSION

On the basis of one year experimentation results concluded that soybean fertilized with 40 kg P₂O₅ ha⁻¹ in combination with 30 kg sulphur ha⁻¹ and foliar spray of GA₃ @ 75 ppm was found beneficial for improving productivity of soybean. These results are only suggestive and require further experimentation to arrive at more consistent and final conclusion.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Chouhan GS. Soybean [*Glycine max* (L.) Merrill] products and their export potential. Green Farming. 2007;1(3):55-7.
2. DAC. & F.W. New Delhi: Directorate of Economics and Statistics, Govt. of India; 2019-20;75.
3. DOA. Rajasthan Agricultural Statistics at a Glance Commissionerate of Agriculture. Jaipur, Rajasthan; 2019-20.

4. Usherwood NR. Nutrient Management for Top-Profit Soybeans; 1998. Available: <http://www.ppi-ppic.org>. (verified 2/15/14)
5. Kanwar JS, Mudahar MS. Fertilizer sulphur and food production. Dordrecht: Martinus-Nijhoff/Dr W. Junk Publisher. 1986;247.
6. Schnug E, Haneklaus E, Booth E, Walker KC. Sulphur and stress resistance in oilseed rape. Proceeding of the 9th International Rapeseed congress. 1995;1(4):229-31.
7. Fazli IS, Abdin MZ, Jamal A, Ahmad S. Interactive effect of sulphur and nitrogen on lipid accumulation, acetyl-CoA concentration and acetyl-CoA carboxylase activity in developing seeds of oilseed crops (*Brassica campestris* L. and *Eruca sativa* Mill.). Plant Sci. 2005;168(1):29-36. DOI: 10.1016/j.plantsci.2004.07.008
8. Solomani A, Sivakumar C, Anbumani S, Suresh T, Arumugam K. Role of plant growth regulators on rice production: A review. Agric Rev. 2001;23:33-40.
9. Aftab T, Masroor M, Khan A, Idrees M, Naeem M, Moinuddin. Salicylic acid acts as potent enhancer of growth, photosynthesis and artemisinin production in *Artemisia annua* L. J Crop Sci Biotechnol. 2010;13(3):183-8. DOI: 10.1007/s12892-010-0040-3.
10. Gomez KA, Gomez VA. Statistical procedure for agriculture research. New York: John Wiley & Sons Inc; 1984.
11. Singh D, Raghuvanshi K, Pandey SK, George PJ. Effect of bio fertilizers on growth and yield of pearl millet (*Pennisetum glaucum* L.). Res. on Environ. Life Sci. 2016;9(3):385-6.
12. Miranda RDS, Ruppenthal V, Lopes LS, Vieira CF, Marques VB, Lacerda F. Phosphorus fertilization improves soybean growth under salt stress. Int. J Plant Anim Sci. 2013;1:21-9.
13. Singh G, Sekhon HS, Ram H, Sharma P. Effect of farmyard manure, phosphorus and phosphate solubilizing bacteria on nodulation, growth and yield of Kabuli chickpea. J Food Legumes. 2010;23:226-9.
14. Chavan PG, Shinde VS, Kote GM, Solunke PS, Bhondve AA. Response of sources and levels of phosphorus with and without PBS inoculation on growth, yield and quality of soybean. Res Crops. 2008;9:286-9.
15. Meena LR, Singh RK, Gautam RC. Effect of moisture conservation practices, phosphorus levels and bacterial inoculation on growth, yield and economics of chickpea (*Cicer arietinum* L.). Legume Res. 2006;29(1):68-72.
16. Kumar V, Chauhan RS, Yadav AS, Upadhyay MK, Rajput RK. Effect of *rhizobium* and phosphorus application on growth, yield and economic of chickpea (*Cicer arietinum* L.). Adv Plant Sci. 2007;20(2):89-90.
17. Singh D, Raghuvanshi K, Chaurasiya A, Dutta SK. Bio fertilizers: nonchemical source for enhancing the performance of pearl millet crop (*Pennisetum glaucum* L.). Bull. Env. Pharmacol. Life Sci. 2017;6(11):38-42.
18. Watimongla J, Gohain T. Effect of phosphorus and sulphur levels on yield, quality of soybean [*Glycine max* (L.) Merrill] and its residual effect on french bean research (*Phaseolus vulgaris*). J Interacademia. 2012;16(2):265-73.
19. Ganeshamurthy AN. Effect of sulphur application on seed and oil yield of soybean. J Ind Soc Soil Sci. 1996;44(2):290-4.
20. Meena DS, Ram B, Tatarwal JP. Productivity, quality and profitability of soybean (*Glycine max* L.) as influenced by sulphur and boron nutrition. Soybean Res. 2011;9:103-8.
21. Vyas AK, Billore SD, Joshi OP, Pachlania N. Productivity of soybean (*Glycine max* L.) genotypes as influenced by nitrogen and sulphur nutrition. Ind J Agric Sci. 2006;76(4):272-83.
22. Sharma OP, Singh GD. Effect of sulphur in conjunction with growth substances on productivity of cluster bean and their residual effect on barley. Ind J Agron. 2005;50(1):16-28.
23. Singh YP, Aggarwal RL. Effect of sulphur levels on yield, nutrient uptake and quality of black gram (*Phaseolus mungo*). Ind J Agron. 1998;43:448-52.
24. Akter A, Ali E, Islam MMZ, Karim R, Razzaque AHM. Effect of gibberellic acid on growth and yield of mustard. Int. J. of Sustai. Crops Prod. 2007;2(2):16-20.
25. Emongor V. Gibberellic acid influence on vegetative growth, nodulation and yield of cowpea (*Vigna unguiculata* (L.) Walp. J Agron. 2007;6:509-17.
26. Noor F. Physiological, biochemical and yield attributes of french bean (*Phaseolus*

- vulgaris* L.) in response to the application of growth regulators [Ph.D. thesis]. Deptt. of Botany. Savar, Dhaka, Bangladesh: Jahangirnagar University. 2014;285.
27. Sapkal KD, Marawar MW, Amarshettiwar SB. Foliar application of plant growth regulators on physiological parameters of soybean. Ann. of Plant Phys. 2011;25(2): 134-6.
28. Vikaria GB, Talpada MM, Suttaria GS. Effect of foliar nutrition of potassium nitrate on the growth and yield of green gram. Legume Res. 2013;36(2): 162-4.
29. Anandha KS, Palchamy A, Mahendran S. Effect of foliar spray of nutrients on growth and yield of green gram. Legume Res. 2004:149-50.

© 2022 Ghasil et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96493>