



Compatibility of Bio and Natural Compounds with *Trichogramma evanescens* (Westwood) for *Pieris rapae* Management in Cauliflower

Amany Nady Siam^a and M. Y. El-Kholy^{b*}

^a Department of Cutworms and Mole Crickets, Plant Protection Research Institute, Agricultural Research Center, Cairo, Egypt.

^b Department of Pests and Plant Protection, National Research Center, Dokki, Cairo, Egypt.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRAF/2023/v9i3216

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100788>

Original Research Article

Received: 19/04/2023

Accepted: 21/06/2023

Published: 09/07/2023

ABSTRACT

The efficacy of a bio compound, Protecto (*Bacillus thuringiensis*) and botanicals extract (mixture of ginger, garlic, hot red pepper extracts and aloe vera) either in combination with *Trichogramma evanescens* releases or alone was evaluated in comparison to an insecticide, Uphold 36% SC (Methoxyfenozide/ Spinetoram) on *Pieris rapae* on cauliflower *Brassica oleracea*. Field experiments were conducted at Atfih District, Giza Governorate, Egypt in September 2020 and October 2021. Results revealed that, all the tested compounds combined with *Trichogramma* releases were more effective in reducing *P. rapae* population than when used alone. *B. thuringiensis* with *Trichogramma* releases caused 96.91% and 95.32% reduction in larval population in 2020 and 2021 respectively while the reduction in *Pieris* pupal pupation was 70.87% and 71.59% respectively during these two trials. Similar results were observed with botanical extracts. Parasitization by *T.*

*Corresponding author: E-mail: melkholy2000@yahoo.com;

evanescens ranged between 91.95-92.77% in the plots sprayed with bio- and natural compounds during the trials in two years. Meanwhile, the insecticide treatment was the least effective in reducing *P. rapae* population (37.85% and 47.85% respectively). It could be concluded that either *B. thuringiensis* or the botanicals extract could be integrated with *T. evanescens* releases for the sustainable production of Cauliflower.

Keywords: *Compatibility; Trichogramma evanescens; Bacillus thuringiensis; botanicals extract; cauliflower; Pieris rapae; sustainable production.*

1. INTRODUCTION

Agriculture faces several challenges like rapid exploitation of the natural resources, the huge use of insecticides and the sudden changes in the global climate, in addition to the need for the production of safe, pesticides-free agricultural products mainly the edible vegetables [1,2]. All these challenges push researchers and those responsible for public health to work on new, safe, available and eco-friendly alternatives to the synthetic pesticides for suppressing pests invading agricultural crops [3]. One of the important vegetable crops is the cauliflower (*Brassica oleracea*). This economically important crop is seriously attacked by the cabbage white butterfly *Pieris rapae* L. (Family: Pieridae). The pest is the most abundant during all the seasons, where the female lays solitary, yellowish eggs, hatching into nasty, green larvae which considered the most serious and destructive stage causing great economic loss in the yield [4] and [5,6]. Continuous application of broad spectrum synthetic insecticides has resulted in the development of the pest resistance, destruction of beneficial insects and contamination for the environment [7]. Currently, great attention has been drawn to the use of various biological agents such as the parasitoids and entomopathogens instead of synthetic insecticides [8], [9], [10] and [11]. *Trichogramma* parasitoids are considered the most efficient and promising organisms in biological control programs [12]. As the damage caused by *P. rapae* reduces the crop production, the intervention with *Trichogramma* parasitoids with compatible bio and natural compounds is imperative to conserve naturally occurring bioagents and to harvest healthy and safe produce [13,14,15]. The entomopathogen *Bacillus thuringiensis* is considered as an efficient biological control agent resulting in significant reduction in the pest and in the use of chemical insecticides [16]. The components of a botanical extract [17] such as ginger, garlic, and hot red pepper, and aloe vera possess an insecticidal effect against many insects [18], [19],

[20] and [21]. The botanicals mixture was found to be effective against certain pests [17] and was safe to *Trichogramma evanescens* Westwood under laboratory conditions [22]. The integration of the parasitoids with bio and natural compounds was earlier evaluated by Oatman et al. [23] and Losey et al. [24] who found that, *T. evanescens* with *B. thuringiensis* was effective against *Helicoverpa zea* Boddie and *Ostrinia nubilalis* (Hubner) . Because of the legal restrictions on the edible fresh vegetables and the periodical sprays of different insecticides, alternatives have to be introduced to suppress *P. rapae* [25,26]. Therefore, this work was carried out to evaluate the efficacy of *B. thuringiensis* , the botanicals mixture extract and combinations with *T. evanescens* releases against *P. rapae* infesting cauliflower crop.

2. MATERIALS AND METHODS

Experiments were carried out to evaluate the integration of *T. evanescens* Westwood releases with certain bio and natural compounds for the management of *P. rapae* on cauliflower. Comparisons were made with an insecticide. Preparation of the botanicals extract and the rearing processes were carried out at the *Trichogramma* Mass Rearing Laboratory of Fayoum Governorate, Plant Protection Research Institute, Agricultural Research Center, Egypt.

2.1 Rearing *T. evanescens*

The rearing of the host *Sitotroga cerealella* and the parasitoid *T. evanescens* were conducted at 25±2 °C and 70±5% R.H. The host *S. cerealella* Olivier was reared as the modification of Hassan [27]. The parasitoid was multiplied on *S. cerealella* eggs.

2.2 Preparation of the Botanicals Extract

The botanicals extract was prepared as per the methods described by Siam and Othman [17]. About one kilogram of garlic bulbs, dry hot pepper, fresh ginger rhizomes were cleaned,

ground well, soaked in ten liters water for 48-72 h and filtered. About one kilogram of aloe vera leaves were cut into small pieces, blended and added to the filtrate. This mixture was diluted to 100 liters water for spraying.

2.3 Field Experiments

Field experiments were carried out at Atfih District, Giza Governorate, Egypt in cauliflower seasons of 2020 and 2021. The experiments carried out as completely randomized block design with six treatments and the control with three replications. Each replicate was ¾ feddan and each plot was separated with three rows as a buffer zone from the adjacent plot to reduce interference of the treatments. The experimental plots were planted with cauliflower seedlings (Colonia variety) on 25th September. Each plot had 10 rows of cauliflower plants which was 50 cm a side within the rows. Plants were inspected for the presence of eggs, larvae and pupae of *P. rapae* every 10 days until harvest on 5th December. Pre-treatment population count was made randomly from half of the rows. Plants had eggs of *P. rapae* was marked with coloured paper cards. Larvae and pupae were also counted. The first release of *T. evanescens* was done after 20 days of transplanting the seedlings. Hard paper envelopes housing the parasitoids were hung in the canopy of the cauliflower plants at the rate of 60 envelopes/ feddan (about 90000 parasitoids/each release). Each envelope had three pieces of different developmental stages of *T. evanescens*. The envelopes were hung every 10 days at the sunset. Normal cultivation practices were followed in cauliflower field.

The treatments were: Releasing *T. evanescens* alone, spraying *B. thuringiensis* alone, spraying the botanicals extract alone, combination of *T. evanescens* with *B. thuringiensis*, combination of *T. evanescens* with the botanicals extract, the chemical insecticide Uphold 36% SC and the control (without any treatment). Back motor sprayer (Agranomodo- China) of 20 liters was used for spraying the compounds. Unparasitized and parasitized *Pieris* eggs, number of both larvae and pupae of *P. rapae* were the criteria used for the evaluation of the treatments. Upon appearance of *Pieris* larvae, sprays of the tested compounds were applied in the early morning with the recommended rates as following:

- 1) *Bacillus thuringiensis* sub sp *kurstaki* 9.4% WP at the rate of 300 gm/200 L. (Product

of Plant Protection Research Institute Agricultural Research Center, Egypt.)

- 2) The prepared botanicals extract at the rate of 1L/100L.
- 3) Uphold 36% SC at the rate 250cm/200L. (Commonly used by farmers in the neighboring fields in case of heavy infestation with the pest)

2.4 Data Collection and Statistical Analysis

The *Pieris* eggs collected from the plots were observed every 10 days to record the parasitism (the black color of the eggs). The efficacy of the treatments was assessed through the percent reduction of *Pieris* population. It was calculated according to Henderson and Tilton [28].

$$\text{Reduction percentage} = \left[1 - \left(\frac{\text{Population of pest after treatment}}{\text{Population of pest before treatment}} \times \frac{\text{Population in control before treatment}}{\text{Population in control after treatment}} \right) \right] \times 100$$

Obtained data of treated plots were compared with those of Uphold insecticide and control plots. Statistical analysis was carried out by IBM-SPSS for windows (v.25). Variance Analysis (ANOVA) was used to process the obtained data. Duncan's multiple range test [29] was used for separation of means [30].

3. RESULTS

3.1 Efficiency of Tested Compounds on *P. rapae*

3.1.1 Reduction percentages of *P. rapae* larvae

The bio compound *B. thuringiensis* combined with *T. evanescens* releases was significantly more effective in reducing *P. rapae* larvae population in 2020 and 2021 (P<0.05) with reduction percentages of 96.91% and 95.32% respectively. This was followed by the natural compound the botanicals extract combined with the parasitoids releases causing the reduction in *Pieris* larvae population by 96.31% and 94.89% in 2020 and 2021 respectively. The chemical insecticide Uphold was the least effective compound in reducing the larvae population with respective reductions of 37.85% and 47.85% (Fig. 1, 3).




Fig. 1. Population of *P. rapae* larvae on cauliflower in 2020 growing season




Fig. 2. Population of of *P. rapae* pupae on cauliflower in 2020 growing season

3.1.2 Reduction percentages of *P. rapae* pupae

The tested compounds significantly differed in its effectiveness against the population density of *P. rapae* pupae ($P < 0.05$). The botanicals extract accompanied with *T. evanescens* releases was effective in reducing pupae of *P. rapae* on cauliflower plants in 2020 and 2021 growing

seasons with reduction percentages of 77.4% and 72.63% respectively, followed by the treatment of combination of *B. thuringiensis* with *Trichogramma* releases with reduction of 70.87% and 71.59% in 2020 and 2021 respectively. The least reductions of 26.12% and 14.04% in *P. rapae* pupal population in 2020 and 2021 respectively were recorded in plots sprayed with the chemical insecticide Uphold. (Fig 2, 4).




Fig. 3. Population of P. rapae Larvae on Cauliflower in 2021 growing season




Fig. 4. Population of P. rapae pupae on Cauliflower in 2021 growing season

3.1.3 Parasitism percentage

There was no parasitism rate recorded before any treatment in 2020 and 2021. With the successive releases of *T. evanescens*, increase in the parasitism was observed in parasitoid-released plots. Results presented in Tables 1, 2 revealed that the plots treated with *T. evanescens* and *B. thuringiensis* had the parasitism rates of 92.56% (2020) and 92.17%

(2021). Also, *T. evanescens* releases with the botanicals extract recorded parasitism rates of 91.95% (2020) and 92.77% (2021). Results revealed that releases of *Trichogramma* parasitoids alone without any compound recorded parasitism rates of 91.03% (2020) and 89.99% (2021). The remaining treatments plots showed no parasitism rate as there no released parasitoids (Tables 1, 2) proving no natural parasitism.

Table 1. Parasitism by *T. evanescens* on *P. rapae* eggs (2020)

<i>T.evanescens</i>			<i>B.thuringiensis</i>			Botanicals extract			<i>T.evanescens</i> + Bt			<i>T.evanescens</i> +botanicals extract			Insecticide			Control		
unp.	P.	%Paras	unp.	P.	%Paras	unp.	P.	%Paras	unp.	P.	%Paras	unp.	P.	%Paras	unp.	P.	%Paras	unp.	P.	%Paras
7.66	0	0	8.33	-	-	8	0	0	8.67	0	-	8.33	0	0	8	-	-	7.66	-	-
3	8.66	74.27	5.67	0	0	5.66	0	0	2.33	9	79.44	2.33	9	79.44	8.67	0	0	9.33	0	0
2	12.33	86.04	6.33	0	0	6.67	0	0	2	12.33	86.04	2.33	11.33	82.94	9.33	0	0	10.33	0	0
0.67	12.33	94.85	5.67	0	0	6	0	0	0.33	12	97.32	0.33	12.33	97.39	9.33	0	0	10.67	0	0
0	12.33	100	6	0	0	6	0	0	0	13	100	0	13	100	9	0	0	13.33	0	0
0	13.33	100	2	0	0	1.67	0	0	0	13	100	0	13	100	9.33	0	0	12.67	0	0
2.22	9.83	91.03	5.67	0	0	5.67	0	0	2.22	11.87	92.56	2.22	11.73	91.95	8.94	0	0	10.67	0	0

Table 2. Parasitism by *T. evanescens* on *P. rapae* eggs (2021)

<i>T.evanescens</i>			<i>B.thuringiensis</i>			Botanicals extract			<i>T.evanescens</i> +Bt			<i>T.evanescens</i> +botanicals extract			Insecticide			Control		
unp.	P.	%Paras	unp.	P.	%Paras	unp.	P.	%Paras	unp.	P.	%Paras	unp.	P.	%Paras	unp.	P.	%Paras	unp.	P.	%Paras
8.33	-	-	7.67	-	-	8	0	0	8.33	-	-	7.67	-	-	7.67	-	-	7.33	-	-
3.33	7.67	69.73	7.33	0	0	7.67	0	0	2.67	8.33	75.73	2.33	8.67	78.82	8.33	0	0	9.67	0	0
2.67	13.11	83.08	7.11	0	0	8.33	0	0	2.03	13.33	86.78	2	13.67	87.24	9.67	0	0	11.33	0	0
0.33	15.33	97.89	6.67	0	0	5.67	0	0	0.23	13.67	98.35	0.33	14.67	97.8	11.33	0	0	15.03	0	0
0	15.76	100	7.33	0	0	6.67	0	0	0	15.33	100	0	15.67	100	13	0	0	18	0	0
0	16.33	100	7.33	0	0	7.67	0	0	0	16.67	100	0	16.33	100	13.67	0	0	19	0	0
2.44	13.62	90.14	7.24	0	0	7.34	0	0	2.21	13.47	92.17	2.01	13.8	92.77	10.61	0	0	13.39	0	0

4. DISCUSSION

The impact of integration between synthetic or natural compounds with bio control agents like *Trichogramma* parasitoids on the latter should be considered. The present work in 2020 and 2021 growing seasons indicates that incorporating either *B. thuringiensis* or the botanicals mixture extract might be compatible with *T. evanescens* for suppressing *P. rapae* populations infesting cauliflower plantations. Obtained results revealed that releasing *T. evanescens* with *B. thuringiensis* could suppress *P. rapae* larvae and pupae populations and resulted in the highest parasitism rates of *Pieris* eggs than those treated by releasing *Trichogramma* parasitoids alone, that's because *Trichogramma* females parasitizes fresh eggs of the pest and *B. thuringiensis* kills larvae of the pest. Our results are in concurrence with those of Lundgren et al. [31] who evaluated releasing *Trichogramma* parasitoids with and without *B. thuringiensis*; their results confirmed that releasing only *Trichogramma* parasitoids was less effective than releasing them with *B. thuringiensis* for controlling Lepidopteron pests. Also, Suzanne et al. [32] supported our observations with their work on controlling *P. rapae* in organic cabbage crop. They reported that *B. thuringiensis* with *Trichogramma* releases was the most effective treatment as it caused high parasitism rate of *Pieris* eggs with a remarkable reduction in larvae populations which resulting in a significant increase in the crop yield over the treatment with *Trichogramma* alone or controls. Other authors worked on the combined use of different species of *Trichogramma* parasitoids and *B. thuringiensis* on different crops aiming at the control of different Lepidopteron pests. Losey et al. [24] proved high efficiency of the combination of *T. nubilalis* and *Bt.*, for controlling *Ostirnia nubilalis* H. (Lepidoptera: Pyralidae) infesting sweet corn. The combination of *T. evanescens* with Agerin (*B. thuringiensis*) was highly effective against *Cryptoblabes gnidiella* Mill infesting grape orchards as it caused the highest parasitism than those with the parasitoids only [33]. AlSaedi et al. [34] reported that *B. thuringiensis* was not harmful for the parasitoids as proven by the highest parasitism rates of two species of *Trichogramma* parasitoids on *Tuta absoluta* infesting tomato in greenhouse. *T. evanescens* was highly effective on suppressing the *Cydia pomonella* (L.) in apple plantations when the parasitoid releases were combined with *B. thuringiensis* than either of the agents being used alone [35-37]. The results from the current

study revealed that *B. thuringiensis* had no adverse effect on *Trichogramma*. Earlier studies agreed with our findings as those of Hwang et al. [38] who stated that, the combination of *Trichogramma ostriniae* (Peng and Chen) and *Bt* reduced the pest population significantly over either of the bio agent alone [25,26]. Ksentini et al. [39] found that among *B. thuringiensis*, deltamethrin and Spinosad, only *Bt* was harmless to the immature developmental stages of *T. cacoeciae* Marchal, *T. bourarache* Pintureau and *T. evanescens* Westwood in pomegranate orchard. Spore mixture of *B. thuringiensis* did not affect the parasitization performance or longevity of the *T. evanescens* adults [40]. Similarly, [41] found that *B. thuringiensis* was safe to *T. evanescens* and was able to suppress *T. absoluta* infesting tomato fields at Fayoum Governorate, Egypt by releasing *T. evanescens* and spraying *B. thuringiensis*. The present study declared that, the use of *B. thuringiensis* as alone recoded reduction in *Pieris* larvae and pupae populations. Our findings are consistent with an earlier studies by Varma et al. [42], Khan et al. [43], Furlong et al. [44] and Maxwell and Fadamiro [45] who reported that *B. thuringiensis* was significantly effective against *Pieris* larvae while El-Fakharany and Hendawy [46] reported that Bio Guard (*B. thuringiensis*) was the least effective compound against *P. rapae*.

In this work, the combination of the botanicals extract with *T. evanescens* releases did not affect the parasitism despite its lethal effect on *Pieris* larvae. Siam and El-Genaidy [22] reported that the botanicals extract was safe to the emergence, fecundity and the general productivity of the *T. evanescens*. The same botanicals extract had a bio pesticide property [17], effective against mango scale insect and was recommended as safe, easy prepared extract and an alternative to synthetic insecticides.

5. CONCLUSION

Either *B. thuringiensis* or the botanicals extract in combination with *T. evanescens* releases has the potential to suppress *P. rapae* populations on cauliflower plants and could be used as alternative for synthetic insecticides.

COMPETING INTERESTS

Authors have declared that they have no known competing financial interests or non-financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. Abd EL Hafez A. A comparison of thermal requirements and some biological aspects of *Trichogramma evanescens* (Westwood) and *Trichogrammatoidea bactrae* (Najgaja) reared from eggs of the pink and spiny bollworm. Ann. Agric. Sci. Ain Shams Univ. Cairo. 1995;4(2):901-912.
2. Embaby ME, Lotfy DE. Ecological studies on cabbage pests. Int. J. Agric. Technol. 2015;11(5):1145-1160.
3. Rani AT, Kammar V, Rani V, Suzan M, Pandey KK, Singh J. Bio pesticides: An alternative to synthetic insecticides. Microbial Technology for Sustainable Environment. 2021:439-466.
4. Ikeura H, Fumiyuki K, Yasuyoshi H. How do *Pieris rapae* search for Brassicaceae host plants? Biochemi. Cal. Systematics and Ecology. 2010;38:1199-1203.
5. Barakat H, Abdul Rasheed W, Douglas GP. Mapping foliage damage index and monitoring of *Pieris brassicae* by using GIS-GPS technology on Cole crops. Journal of Entomology and Zoology Studies. 2018;6(2):933-938.
6. ELSayed EI. Neem *A. indica* seeds as anti-feedant and ovi-positional repellent for the Egyptian cotton leafworm *Spodoptera littoralis* (Biosd.). Bull. Ent. Soc. Egypt. Econ. Ser. 1982;(49):49-58.
7. Delobel A, Malonga P. Insecticidal properties of six plant materials against *Caryedon serratus* Oliv. J. Stored Prod. Res. 1987;23(3):173-176.
8. Salem SA, Abdel-Rahman AA, Zebitz CPW, Saleh MME, Ali IF, El-Khouly MY. Evaluation of entomopathogenic nematodes in controlling some cabbage pests. J.Appl.Sci.Res. 2007;3(4):323-328.
9. Gutierrez CG, Garcia NM, Campus MN, Hernandez IC. Efficacy of *Beauveria bassiana* and *Metarhizium anisoplae* to control *Pieris rapae* on cabbage in the field. Southwestern Entomologist. 2010;35(1):75-83.
10. Gosh SK. Harmful effect of insecticides in the population dynamics of spiders on ladies'finger *Abelmoschus esculentus* (L.) Moench at field level. Am- Euras. J. Agric. & Environ. Sci. 2013;13(9):1181-1186.
11. Halder J, Seni A. Biological management of major vegetable insect pests with macro and microorganisms. Microbes for Sustainable Insect Pest Management. 2021:233-252.
12. Zang LS, Wang S, Zhang F, Desneux N. Biological control with *Trichogramma* in China: History, Present status and perspectives. Annual Reviews of Entomology. 2021;66(24):1-22.
13. Golob O, Mwumbola JV, Ngulube F. The use of locally available materials as protectants of maize grains against insects' infestation during storage in Malawi. J. Stored. Prod. Res. 1982;18:67-74.
14. Goncalves EM, Alegria C, Abreu M. Benefits of *Brassica* nutraceutical compounds on human health. Characterization, Functional Genomics and Health benefits. 2013;19:2013
15. Krishnamoorthy A. Exploitation of egg parasitoids for control of potential pests in vegetable ecosystems in India. Comunicata Scientiae. 2012;3:1-5.
16. Alejandra B, Supaporn L, Sarjeet SG, Mario S. *Bacillus thuringiensis*: A story of a successful bio insecticide. Insect Biochemistry and Molecular Biology. 2011;41(7):423-431.
17. Siam A, Othman E. Field evaluation of botanicals extracts for suppressing the mango scale insect, *Aulacaspis tubercularis* (Newstead) (Hemiptera: Diaspididae). Egyptian Journal of Biological Pest Control. 2020;30(1):1-5.
18. Okonkwo CO, Ohaeri OC, Insecticidal potentials of some selected plants. J. of Chem. and Pharma. Resear. 2013;5(4):370-376.
19. Teklay M, Prasad SHKR, Etana B, et al. Insecticidal and repellent properties of selected medicinal plants collected from Sofoho, Axum, North East Africa. Int.J.Sci.Inn.Tech.Sec.A. 2012;1(3):1-8.
20. Neasel A, Charleston BG, et al. Testing the insecticidal potential of Chili pepper (*Capsicum frutescens*) fruit extract against termites (*Coptotermes gestroi*). Bio. 182. Biotech; 2016.
21. Karkanis CA, Christos GA. Natural insecticides from native plants of Mediterranean basin and their activity for the control of major insect pests in vegetable crops: Shifting from the past to the future. Journal of Pest Science. 2021;94(2):187-202.
22. Siam A, El-Genaidy M. Laboratory evaluation of certain natural plants extracts on the fitness components of *Trichogramma evanescens* (Westwood) (Hymenoptera: Trichogrammatidae). Bioscience Research. 2021;18(1):739-748.

23. Oatman ER, Wyman JA, Van Steenwyk RA, Johnson MW. Integrated control of the tomato fruit worm (Lepidoptera: Noctuidae) and other Lepidopterous pests on fresh-market tomatoes in Southern California. *Journal of Economic Entomology*. 1983;76:1363-1369.
24. Losey JE, Fleischer SJ, et al. Evaluation of *Trichogramma nubilalis* and *Bacillus thuringiensis* in management of *Ostrinia nubilalis* (Lepidoptera: Pyralidae) in sweet corn. *Environmental Entomology*. 1995;24:436-445.
25. Knipling EF, Mc Gure JU. Population models to appraise the limitations and potentialities of *Trichogramma* in managing host insect population. *USDA Technical Bulletin*. 1968;1387.
26. Nancy ME, Wendy MC. Alternatives to synthetic chemical insecticides for use in crucifer crops. *Biological Agriculture & Horticulture*. 2012;8(1):33-52.
27. Hassan SA. Improved method for the production of the Angoumois grain moth *Sitotroga cerealella* (Oliv.). *Trichogramma* and other egg parasitoids. Cairo, Egypt. 1995;(4-7):157-170.
28. Henderson CF, Tilton EW, Tests with Acaricides against the brown wheat mite. *J. Econ. Entomol.* 1955;48(2):157-161.
29. Duncan DB. Multiple range and multiple F tests. *Biometrics*. 1955;11:1-41.
30. Snedecor GW, Cochran WG. *Statistical methods*. 2nd The Iowa State University Press Ames Iowa; 1980.
31. Lundgren LG, Heimpel GE, Bomgren SA. Comparison of *Trichogramma brassicae* (Hymenoptera: Trichogrammatidae) augmentation with organic and synthetic pesticides for control of cruciferous Lepidoptera. *Environmental Entomology*. 2002;31:1231-1239.
32. Suzanne B, Rayan A, Adams S, Franklin J, Driscoll J. Use of *Trichogramma* and *Bt* for control of *Pieris rapae* in cabbage on Prince Edward Island. *J. Acad. Entomol. Soc.* 2015;11:26-31.
33. Tohamy TH, et al. Biological control of the Honey Dew Moth *Cryptoblabes gnidiella* Mill in grape orchards using the local egg parasitoid *Trichogramma evanescens* (Westwood) and Agerin (*Bacillus thuringiensis*) compared with recommended insecticides in Middle Egypt. *Assiut.J.of Agric.Sci.* 2007;38(3):93-111.
34. AlSaedi G, Ashouri A, Reza TH. Assessment of two *Trichogramma* species with *Bacillus thuringiensis* var. *Kurstaki* for the control of tomato leaf miner *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) in Iran. *Open Journal of Ecology*. 2017;(7):112-124.
35. Oztemiz S, Kuden A, Serpil N, Lavkor I. Efficacy of *Trichogramma evanescens* and *Bacillus thuringiensis* var. *kurstaki* in control of *Cydia pomonella* (L.) in Turkey. *Turkish Journal of Agriculture and Forestry*. 2017;41(3):201-207.
36. Picchi V, Fibiani M, Scalzo LR. Chapter 2. Cauliflower. Nutritional composition and antioxidant properties of fruits and vegetables. 2020:19-32. Available:<http://doi.org/10.1016/B978-0-12-812780-3-00002-7>
37. Pourarian S, Shirazi J, Rasipour A. An investigation on the biology and efficiency of *Trichogramma* spp. on the eggs of *Pieris rapae* under laboratory conditions. *Bio control in Plant Protection*. 2017;4(2):39-53.
38. Hwang IC, Park C, Kang DK, Jin NY. Combined application of *Trichogramma ostriniae* and *Bacillus thuringiensis* for eco-friendly control of *Plutella xylostella*. *Journal of the Korean Society for Applied Biological Chemistry*. 2010;53(3).
39. Ksentini I, Jardak T, Zeghal N. *Bacillus thuringiensis*, deltamethrin and spinosad side effects on three *Trichogramma* species. *Bulletin of Insectology*. 2010;63(1):31-37.
40. Azizoglu U, Semih Y, Ayvaz A, Salih K. Effect of *Bacillus thuringiensis* sub. *kurstaki* HD1 spore-crystal mixture on the adults of egg parasitoid *Trichogramma evanescens* (Hymenoptera: Trichogrammatidae). *Agriculture and Environmental Biotechnology*. 2015;29(4):653-658.
41. Siam A. Studies on some biological and ecological factors affecting the performance of *Trichogramma evanescens* (Westwood) under laboratory and field conditions. Ph. D. thesis. Cairo University, Egypt; 2017.
42. Varma GC, Bindrs OS, Singh D, Darshan S. Comparative efficacy of *Bacillus thuringiensis* Berliner formulations against *Pieris brassicae* Linnaeus. *Current Science*, Department of Entomology, Punjab University, Ludhiana. India. 1974;43:734-735.
43. Khan HH, Kumar A, Naz H. Evaluation of chemical and biological insecticides for the management of cabbage butterfly *Pieris rapae* (Linn.)(Lepidoptera: Pieridae)- A review. *Journal of Pharmacognosy and phytochemistry*. 2017;6(6):2224-2233.

44. Furlong MJ, Ju KH, Su PW, Chol KJ, Chang II, Zalucki P. Integration of endemic natural enemies and *Bacillus thuringiensis* to manage insect pests of *Brassica* crops in North Korea. *Agriculture, Ecosystems & Environment*. 2008;125(1-4):223-238.
45. Maxwell EM, Fadamiro HY. Evaluation of several reduced-risk insecticides in combination with an action threshold for managing lepidopteran pests of Cole crops in Alabama. *Florida Entomologist*. 2006;89(2):117-126.
46. El-Fakharany SKM, Hendawy AS. Field studies on cabbage white butterfly *Pieris rapae* (Linnaeus) and its associated parasitoid and predatory species in Egypt. *Egyptian Journal of Biological Pest Control*. 2014;24(2):437-444.

© 2023 Siam and El-Kholy; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/100788>