



## Effect of Establishment Methods and Nutrient Management Practices on Growth Characters and Yield of Rice (*Oryza sativa* L.)

G. Santhosh Kumar <sup>a\*#</sup>, R. Mahender Kumar <sup>b≡</sup>, A. Sreenivas <sup>c<sup>o</sup></sup>, G. Jayasree <sup>d†</sup>  
and T. Ramesh <sup>e‡</sup>

<sup>a</sup> Department of Agronomy, Agricultural College, Warangal -506007, India.

<sup>b</sup> Department of Agronomy, Indian Institute of Rice Research, Rajendranagar, Hyderabad-500030, India.

<sup>c</sup> Department of Agronomy, PJTSAU, Rajendranagar, Hyderabad-500030, India.

<sup>d</sup> Department of SSAC, College of Agriculture, PJTSAU, Rajendranagar, Hyderabad-500030, India.

<sup>e</sup> Department of Plant Physiology, College of Agriculture, PJTSAU, Rajendranagar, Hyderabad-500030, India.

### Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/IJECC/2021/v11i1230657

#### Editor(s):

(1) Dr. Wen-Cheng Liu, National United University, Taiwan.

#### Reviewers:

(1) Ahsan A. Kadhimi, Al-Qadisiyah University, Iraq.

(2) Deivaseeno Dorairaj, Universiti Kebangsaan Malaysia, Malaysia.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/79936>

Original Research Article

Received 22 October 2021

Accepted 25 December 2021

Published 28 December 2021

### ABSTRACT

**Aim:** Study was conducted to compare and to identify the best combination of rice crop establishment method and nutrient management in terms of crop growth, development and yield.

**Study Design:** Experiment was laid out in splitplot design.

**Place and Duration of Study:** The field experiments were conducted at the research farm of Indian Institute of Rice Research, Hyderabad, Telangana State, India during the kharif and rabi seasons of 2011-12 and 2012-13.

# Assistant Professor;

≡ Principal Scientist & Head;

<sup>o</sup> Professor & Univ Head (Retd.);

† Professor;

‡ Professor & Head;

\*Corresponding author: E-mail: santhuugupta@gmail.com;

**Methodology:** The treatments consisted of four establishment methods (System of rice intensification (SRI), modified drum seeder, normal drum seeder and normal transplanting) as main plot treatments and four nutrient management practices (100% recommended dosage of nitrogen (RDN) through inorganic, 75% RDN through inorganic+ 25% RDN through organic, 50% RDN through inorganic+50% RDN through organic and 100% RDN through organic) as sub plot treatments with 16 treatment combinations and three replications. The semi dwarf, high yielding Sampada variety was raised during the study period.

**Results:** Growth parameters like plant height, leaf area index (LAI), dry matter accumulation, number of tillers per square meter were observed at 45,75,105 days after transplanting and at harvest stages. The two years experimental results and the pooled mean data revealed that SRI recorded significantly superior growth parameters at all the crop growth stages followed by modified drum seeder. In respect of subplots 50% RDN through inorganic+50% RDN through organic resulted in superior growth parameters and yield.

**Conclusions:**The combination of SRI with nutrient management practice 50% nitrogen through urea and 50% nitrogen through vermicompost resulted in greater growth parameters and grain yield.

*Keywords: SRI; direct seeding; modified drum seeder; INM.*

## 1. INTRODUCTION

Rice plays a central role in food security and economic growth of India. It is grown in more than one-fifth of the total gross cropped area and contributes more than one-fourth of the total calorie intake [1]. It is the major source of calories for 40% of the world population [2]. About 77% of the global rice production in the world is produced through conventional transplanting methods in puddled soil [3,4]. Conventional transplanting system of rice crop production requires high inputs of labour, water, capital, and energy and hence it has become less profitable at present due to the lack of these resources [3,5]. A shortage of labour during peak periods increase labour wages and make transplanting operation costly [6]. Thus, the farmers prefer direct seeding method as a viable alternative to the drudgery and labour intensive conventional transplanting system.

Direct seeding probably is one of the oldest method of crop establishment which is now practiced in many Asian countries [7]. Direct rice seeding has many advantages to the farmers such as higher economic returns, short crop duration, requires less labour and water suitable for mechanization and have less methane emission [8,9].

Another major concern in rice production systems is the dwindling trend of availability of water resources [10]. Out of 70-80% freshwater used in agriculture, rice accounts for 85% in total and 30% in puddling only. A 10% increase in irrigation efficiency can help bring additional 14 million ha area under irrigation [11]. Hence there is a need to develop and adopt water saving

methods in rice cultivation so that production and productivity levels are elevated despite the looming water crisis [11,12]. System of rice intensification (SRI) method was found to save 22 to 38 per cent of water during dry and wet season, respectively over conventional transplanting of rice establishment [13]. Efficient water management and nutrients could be better utilized under SRI along with integration of nutrient sources to realize the maximum rice crop productivity through enhancement of growth and physiological aspects of lowland rice [14].

Intensive agriculture and decreasing the use of organic manures, have led to severe degradation of soil fertility and productivity of rice cropping systems [15]. The decline or stagnation in yield has been attributed to nutrient mining and reduced use of organics manures [16]. Under high input production systems where productivity cannot be further increased with incremental use of mineral fertilizers alone, the addition of organic sources could increase yield through increased soil productivity and higher fertilizer use efficiency [17,18,19]. Singh and Kumar [20] reported increased yield and nutrient use efficiency in rice with organic manures. The combined use of organic and inorganic fertilizers has been reported to not only meet the nutrient needs of the crop but also has been found to sustain large scale productivity goals [21]. A synergistic effect on crop yield is reported through common use of fertilizer and organics has improved soil fertility. Integrated nutrient management involves maintenance of soil fertility, sustainability of crop production and the beneficial effect of integrated plant nutrient supply (IPNS) in lowland rice has been well reported by several researchers [22].

## 2. MATERIALS AND METHODS

The present investigation was conducted during kharif and rabi seasons of 2011-12 and 2012-13 at Indian Institute of Rice Research farm, Rajendranagar, Hyderabad, Telangana state, India. The farm is geographically situated at an altitude of 542.7 m above mean sea level on 17° 19' N latitude and 78° 29' E longitudes. According to Troll's climatic classification, it falls under semi-arid tropics (SAT). The soil was clay loam, alkaline in reaction (pH 8.0 -8.2), with 0.48-0.52% of organic matter, 210-223 kg ha<sup>-1</sup> of available nitrogen, 39-43 kg ha<sup>-1</sup> of available P and 525-542 kg ha<sup>-1</sup> of available K. The semi dwarf, high yielding rice variety Sampadha, having crop duration of 135 days with yield potential of 5.8-6.8 t ha<sup>-1</sup> was grown in the experimental site. Usually normal drum seeders are available with close spacing (20x5-8 cm<sup>2</sup>), but for this study a new drum seeder was fabricated with 25x25 cm<sup>2</sup> spacing to test the wider spacing efficiency in direct seeding and this treatment was denoted as modified drum seeder. The experiment was laid out in splitplot design with sixteen treatments and three replications. The treatment combinations consisted of four establishment methods i.e., system of rice intensification (SRI) (M<sub>1</sub>), modified drum seeder (25x25cm<sup>2</sup> spacing) (M<sub>2</sub>), normal drum seeder (M<sub>3</sub>) and normal transplanting (NTP) (M<sub>4</sub>) in mainplots and four nutrient management practices i.e., 100% RDN (recommended dose of nitrogen) through inorganic (N<sub>1</sub>), 75% RDN through inorganic+ 25% RDN through organic (N<sub>2</sub>), 50% RDN through inorganic+50% RDN through organic (N<sub>3</sub>) and 100% RDN through organic (N<sub>4</sub>) in sub plots. In case of NTP and SRI the sprouted seeds were broadcasted uniformly in well prepared seed beds. And on the same day seeds were sown in the respective mainfield plots for drum seeder and modified drum seeder. The recommended dose of fertilizer was 120 Kg N: 60 Kg P<sub>2</sub>O<sub>5</sub> : 40 Kg K<sub>2</sub>O ha<sup>-1</sup>. The recommended dose of phosphorus @ 60 kg P<sub>2</sub>O<sub>5</sub> kg ha<sup>-1</sup> through single super phosphate (SSP), potassium @ 40 kg K<sub>2</sub>O ha<sup>-1</sup> as muriate of potash (MOP) and ZnSO<sub>4</sub>@ 20 kg ha<sup>-1</sup> were applied to all the treatments uniformly as basal application. In case of 100% inorganic treatment nitrogen was applied through urea in three equal splits as ½ as basal, ¼ at maximum tillering and ¼ at panicle initiation stage. In INM treatments urea was applied as the inorganic source of nitrogen in three equal splits at the time of transplanting, 30 DAT (days after transplanting) and at 60 DAT. Every season

vermicompost was analysed for nitrogen content and based on nitrogen percentage vermicompost was applied as basal. Twelve days and twenty one days old seedlings were used for transplanting in SRI and NTP respectively. Spacing of 25x25cm<sup>2</sup> was maintained in SRI and modified drum seeder methods and for NTP it was 20x15 cm<sup>2</sup>. Statistical analysis of the data was done as per the standard analysis of variance technique for the experimental designs following INDOSTAT software based programme. The treatment means were compared at P<0.05 level of probability using t test and calculating CD values.

## 3. RESULTS AND DISCUSSION

### 3.1 Plant Height

The experimental results revealed that, the average plant height increased linearly up to 105 DAS and continued to increase until maturity. From the data, it can be inferred that maximum plant height was observed with SRI followed by modified drum seeder at all the crop growth stages over normal drum seeder and NTP (Table 1). Taller plants resulted by adopting SRI method might be due to transplanting of younger and single seedling by keeping the roots straight (ensuring that the roots do not assume 'J' shape). Wider spacing maintenance in SRI and modified drum seeder in both row to row and plant to plant might have encouraged vigorous root system and the plants get sufficient space above the ground to grow and the increased light transmission in the canopy thus leading to greater plant height. Similar results were reported by [23,24,25,8] and [26]. Among the nutrient management treatments, at all the stages, 50% RDN through inorganic+50% RDN through organic treatment recorded significantly higher plant height, followed by 100% RDN through inorganic treatment and 75% RDN through inorganic+25% RDN through organic treatment, these findings are in accordance with [21,15]. The interaction effect was non-significant among planting methods and nutrient management practices in respect of plant height.

### 3.2 Leaf Area Index (LAI)

At all the stages LAI recorded was significantly higher in SRI followed by modified drum seeder over normal drum seeder and NTP during both the years (Table 2). The LAI recorded in modified drum seeder was significantly on par with SRI during kharif 2012-13, rabi 2011-12 and 2012-13.

The mean percentage increases of LAI in SRI at 105 DAS were 48.01, 43.21, 37.70 and 34.01% over NTP during kharif 2011, kharif 2012, rabi 2011-12, rabi 2012-13 respectively (Table 2). This might be due to planting in square geometry with wider spacing and single seedling which facilitated for better utilization of the resources to obtain maximum leaf area and higher number of leaves. These results are in conformity with findings of [26,27,14] and [23].

Maximum LAI was observed with the treatment 50% RDN through inorganic+50% RDN through vermicompost at 105 DAS during all the seasons of study. 100% RDN through vermicompost registered lowest LAI at all the stages of crop growth. The treatment 50% RDN through inorganic+50% RDN through organic influenced the plant growth, tiller number and dry matter accumulation and thus manifested higher LAI. [15,16,27,14] and [24] also expressed the similar opinion.

The interaction effect among planting methods and nutrient management practices on leaf area index was found significant at 45 DAS during kharif 2011 and pooled kharif mean (Table 2a). The interaction was also significant at 75 DAS during kharif 2011, kharif 2012, kharif pooled mean (Table 2a), rabi 2011, rabi pooled mean at 75 DAS in both the years and in pooled means (Table 2b). The data on interaction revealed that at all these stages SRI in combination with 50% RDN through organic+50% RDN through inorganic source registered higher LAI as compared to all other establishment and nutrient management combinations.

### 3.3 Dry Matter Production (kg ha<sup>-1</sup>)

Increases in dry matter production of rice was rather slow up to 45 DAS, thereafter it increased linearly up to 105 DAS and further, it continued to increase slowly until maturity (Table 3).

At 45 DAS the numerically higher values were observed with SRI (1989.8, 2607, 2290.1 and 3126.5 g of total dry matter m<sup>-2</sup> during 2013 and 2014, respectively) followed by modified drum seeder, over other treatments (Table 3). The similar trend was also observed at 75, 105 DAS and at harvest. Highest values of dry matter accumulation was observed at harvest with SRI which was 17.49, 23.57, 14.79 and 23.72% higher as compared to normal transplanting during 2011 kharif, 2012 kharif, 2011-12 rabi and 2012-13 rabi. The higher dry matter production in SRI planting method was

attributed to planting of young and single seedling at shallow depth in wider spacing and two direction cono-weeding which lead to taller plants, higher leaf area, LAI, better root growth, profuse and strong tillers with higher crop growth rate. The results obtained in this investigation are in conformity to the findings of [26,25].

Among all the nutrient management treatments 50% RDN through inorganic+50% RDN through organic recorded significantly higher dry matter accumulation, followed by 100% inorganic treatment, over other treatments. At all the stages of crop growth lowest dry matter accumulation was observed with 100% organic treatment. At harvest 100% organic treatment recorded 16.05%, 9.49%, 13.03%, 9.59%, 11.23% and 12.55% lower dry matter accumulation as compared to 50% inorganic+50% organic in kharif 2011, kharif 2012, rabi 2011-12 and 2012-13, in pooled mean kharif and in pooled mean rabi respectively. These findings were in conformity with the results of [16] and [15].

### 3.4 Number of Tillers m<sup>-2</sup>

The average number of tillers m<sup>-2</sup> of rice increased linearly up to 90 DAS. Narrow spacing in normal drum seeder and normal transplanting, wider spacing in SRI and modified drum seeder led to significant difference in number of tillers m<sup>-2</sup>. But number of tillers m<sup>-2</sup> recorded was significantly higher in SRI over modified drum seeder at 60 and 90 DAS and of pooled means, respectively (Table 4). At all the growth stages NTP recorded significantly lower number of panicles m<sup>-2</sup> as compared to other establishment methods. As compared to normal transplanting SRI practice gets the benefit of the early phyllochron stages (less than four leaves) having higher potential to quick recovery and to produce more tillers. If transplanting is done beyond the fourth phyllochron (after about 15 days), the first primary tiller does not emerge and all of the descendents of this tiller are lost. Similarly, if transplanting is further delayed by the length of another phyllochron, the second primary tiller and all its descendents are also forgone. Planting in square method with wider spacing resulted in profuse tillering under SRI cultivation and facilitated plants for better utilization of the resources. Similar findings have also been reported by [26,8,9,25].

In both the years application of 50% RDN through inorganic source+50% RDN through organic source recorded significantly higher tiller

number  $m^{-2}$  over other all nutrient management practices. Several researchers observed similar results. 50% inorganic fertilizer with 50% organic source improved the general soil environment, physicochemical and biological conditions thus favouring the increased and timely availability of macro and micro nutrients helped in profuse tillering.

### 3.5. Grain Yield ( $kg\ ha^{-1}$ )

The higher grain yield of  $6535\ kg\ ha^{-1}$  &  $6140\ kg\ ha^{-1}$  was recorded by SRI method during 2012 & 2011 kharif seasons respectively. Next to SRI method modified drum seeder proved its significant superiority over normal drum seeder and normal transplanting. During 2011 kharif modified drum seeder remained at par with normal drum seeder but was found significantly superior over normal transplanting normal drum seeder (Table 5). The pooled data also indicated that SRI method stood first with grain yield of  $6337.5\ kg\ ha^{-1}$  followed by modified drum seeder, normal drum seeder and normal transplanting. In terms of percentage increase in yield due to SRI over modified drum seeder, normal drum seeder and normal transplanting was 9.27, 18.24, 21.74% respectively. During rabi season of 2011-12 and 2012-13 SRI method was found significantly superior than the remaining three crop establishment methods. There was high yield difference of  $579\ kg\ ha^{-1}$  and  $358\ kg\ ha^{-1}$  between SRI and modified drum seeder in first and second rabi seasons respectively. The pooled data also showed the advantage of  $468\ kg\ ha^{-1}$  by SRI over modified drum seeder. These results were in accordance with the findings of [28,29].

The yield advantage due to SRI over conventional planting was mainly due to more number of tiller production per unit area accompanied by maximum panicle bearing tillers with low spikelet sterility. Since planting of young seedlings of 12 days in main field with immediate establishment have facilitated early initiation of tillers. It is evident that highest tillers production was observed with SRI planting. Controlled irrigation also augmented the fresh root production till flowering stage and does helped in supplementation of nutrient requires for supporting of filling capacity of panicles.

Among the subplots 50% RDN through inorganic + 50% RDN through organic proved its superiority during all the seasons of experiment. During first kharif season 50% RDN through inorganic + 50% RDN through organic remained at par with 100%

RDN through inorganic. But 2012 kharif data and the kharif pooled data indicated the significant superiority of 50% RDN through inorganic + 50% RDN through organic source. Both the rabi seasons data revealed that 50% RDN through inorganic + 50% RDN through organic was on par with 100% RDN through inorganic and found significantly better over remaining other treatments. The results emphasize the concepts of INM for high grain production and also sustainability of soil fertility [30,31]. During all the seasons 100% organic treatment recorded lowest grain yield [32].

The interaction effect of planting methods and nutrient management practices on grain yield during all the seasons of study and in pooled means was found to be significant (Table 5a). During kharif 2011, kharif 2012, rabi 2011-12, rabi 2012-13 and in kharif and rabi pooled means showed that SRI in combination with 50% RDN through inorganic source + 50% RDN through organic source recorded significantly higher grain yield over other establishment and nutrient combinations.

### 3.6 Straw Yield ( $kg\ ha^{-1}$ )

Straw yield of rice was significantly higher in SRI and during kharif and rabi seasons of 2012-13 it was significantly on par with modified drum seeder treatment (Table 6). During all the seasons straw yield recorded by modified drum seeder was statistically on par with normal drum seeder. By observing the data of all the four seasons it was evident that wider spacing treatments recorded higher straw yield as compared to closer spacing treatments. It was probably due to more dry matter production per unit area caused by better nutrient absorption from soil, increased rate of metabolic processes, higher rate of light absorption and increased rate of photosynthetic activity that produced higher plant height and leaf area index as compared to normal transplanting. These results are in agreement with the findings of [28,29,33].

During all the four seasons and in pooled means highest straw yield was observed with treatment 50% RDN through inorganic source + 50% RDN through organic source. The superiority of this treatment was because of adequate supply of nitrogen throughout crop growth period that led to higher dry matter production [31]. The lowest straw yield was found with treatment 100% organic during all the seasons [32].

**Table 1. Plant height (cm) at different stages of crop growth as influenced by establishment methods and nutrient management practices**

	45 DAS		75 DAS		105 DAS		At harvest	
	Kharif	Rabi	Kharif	Rabi	Kharif	Rabi	Kharif	Rabi
<b>MAIN PLOTS</b>								
M1 (SRI)	37.75	40.90	76.58	82.68	93.49	100.61	101.60	106.30
M2 (Modified Drum Seeder)	34.10	36.10	69.22	70.83	87.00	91.59	90.50	95.50
M3 ( Normal Drum seeder)	34.50	36.50	70.78	80.08	79.61	84.01	89.70	94.70
M4 (NTP)	32.00	32.50	60.57	66.09	74.36	80.73	83.10	89.50
S.E.M	0.40	0.80	1.88	0.97	1.62	1.59	2.30	2.20
CD ( $P=0.05$ )	1.30	2.80	6.49	3.36	5.60	5.49	8.00	7.50
<b>SUBPLOTS</b>								
N1 (100% Inorg)	35.40	37.80	71.45	76.45	85.82	91.18	93.40	98.60
N2 (75% Inorg+25% org)	32.70	34.30	67.00	73.11	80.21	86.23	87.40	94.00
N3 (50% Inorg+50% org)	39.20	41.60	76.82	81.25	93.33	96.15	101.20	104.10
N4 (100%org)	30.90	32.40	61.87	68.86	75.11	83.38	82.90	89.20
S.E.M	0.80	1.00	1.34	1.33	1.68	1.69	1.90	2.00
CD ( $P=0.05$ )	2.40	2.80	3.91	3.89	4.90	4.92	5.60	5.90
MEAN	34.60	36.50	69.28	74.92	83.62	89.23	91.20	96.50
<b>INTERACTIONS</b>								
<b>Sub at same level of Main</b>								
S.E.M	1.70	1.90	2.68	2.66	3.36	3.37	3.90	4.00
CD ( $P=0.05$ )	NS	NS	NS	NS	NS	NS	NS	NS
<b>Main at same level of Sub</b>								
S.E.M	1.50	1.90	2.99	2.50	3.33	3.32	4.10	4.10
CD ( $P=0.05$ )	NS	NS	NS	NS	NS	NS	NS	NS

**Table 2. LAI at different stages of crop growth as influenced by establishment methods and nutrient management practices**

	45 DAS						75 DAS						105 DAS						At harvest					
	Kharif		Rabi		Pooled		Kharif		Rabi		Pooled		Kharif		Rabi		Pooled		Kharif		Rabi		Pooled	
MAIN PLOTS	2011	2012	Pooled	2011	2012	pooled	2011	2012	Pooled	2011	2012	Pooled	2011	2012	Pooled	2011	2012	Pooled	2011	2012	Pooled	2011	2012	Pooled
M1 (SRI)	1.44	1.63	1.54	1.73	2.00	1.86	4.31	3.91	4.11	4.72	5.14	4.93	5.98	5.70	5.84	6.61	7.04	6.82	5.35	5.30	5.32	5.86	6.32	6.09
M2 (Modified Drum seederr)	1.30	1.45	1.37	1.47	1.60	1.54	3.38	3.25	3.31	3.62	4.17	3.90	5.51	5.15	5.33	6.05	6.47	6.26	4.72	4.94	4.83	5.59	5.91	5.75
M3 ( Normal Drum seeder)	1.06	1.31	1.19	1.19	1.38	1.28	2.58	2.36	2.47	3.31	3.71	3.51	4.65	4.57	4.61	5.32	5.96	5.64	4.01	4.35	4.18	4.91	5.31	5.11
M4 (NTP)	0.84	1.15	1.00	1.07	1.19	1.13	2.21	2.24	2.22	2.69	3.00	2.85	4.04	3.98	4.01	4.80	5.25	5.02	3.59	3.72	3.65	4.44	4.87	4.65
S.E.M	0.03	0.03	0.02	0.05	0.06	0.04	0.07	0.07	0.06	0.13	0.15	0.11	0.18	0.13	0.12	0.18	0.21	0.15	0.15	0.16	0.05	0.18	0.20	0.14
CD (P=.05)	0.10	0.11	0.06	0.18	0.20	0.15	0.25	0.25	0.20	0.44	0.51	0.40	0.63	0.46	0.42	0.63	0.74	0.52	0.51	0.57	0.17	0.62	0.68	0.47
SUBPLOTS																								
N1 (100% Inorg)	1.33	1.53	1.43	1.48	1.64	1.56	3.28	3.11	3.19	3.84	4.24	4.04	5.26	5.19	5.23	5.88	6.40	6.14	4.62	4.90	4.76	5.38	5.79	5.58
N2 (75% Inorg+25% org)	1.00	1.25	1.13	1.25	1.44	1.34	2.84	2.63	2.74	3.34	3.75	3.55	4.82	4.62	4.72	5.45	6.01	5.73	4.17	4.32	4.24	4.87	5.41	5.14
N3 (50% Inorg+50% org)	1.52	1.70	1.61	1.63	1.88	1.75	3.98	3.71	3.84	4.22	4.71	4.47	5.71	5.48	5.59	6.40	6.82	6.61	5.10	5.18	5.14	5.91	6.15	6.03
N4 (100%org)	0.79	1.06	0.93	1.09	1.23	1.16	2.39	2.31	2.35	2.93	3.32	3.12	4.39	4.11	4.25	5.05	5.49	5.27	3.78	3.91	3.85	4.63	5.06	4.85
S.E.M	0.03	0.04	0.02	0.05	0.05	0.04	0.08	0.08	0.07	0.13	0.13	0.08	0.17	0.16	0.14	0.18	0.21	0.11	0.16	0.15	0.13	0.18	0.18	0.14
CD (P=.05)	0.08	0.12	0.06	0.13	0.14	0.11	0.23	0.24	0.19	0.37	0.37	0.23	0.49	0.46	0.40	0.53	0.60	0.32	0.48	0.45	0.39	0.53	0.53	0.42
MEAN	1.16	1.39	1.27	1.36	1.54	1.45	3.12	2.94	3.03	3.58	4.00	3.79	5.04	4.85	4.95	5.69	6.18	5.94	4.42	4.58	4.50	5.20	5.60	5.40
INTERACTIONS																								
Sub at same level of Main																								
S.Em±	0.06	0.08	0.04	0.09	0.10	0.07	0.16	0.16	0.13	0.25	0.25	0.15	0.34	0.31	0.27	0.36	0.41	0.22	0.33	0.31	0.27	0.36	0.36	0.29
CD (P=.05)	0.16	NS	0.12	NS	NS	NS	0.47	0.48	0.38	0.74	NS	0.45	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Main at same level of Sub																								
S.Em±	0.06	0.08	0.04	0.09	0.10	0.08	0.16	0.16	0.13	0.25	0.26	0.18	0.34	0.30	0.26	0.36	0.42	0.24	0.32	0.31	0.24	0.36	0.37	0.28
CD(P=.05)	0.17	NS	0.12	NS	NS	NS	0.47	0.49	0.39	0.78	NS	0.55	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

**Table 3. Dry matter accumulation (kg ha<sup>-1</sup>) at different stages of crop growth as influenced by establishment methods and nutrient management practices**

	45 DAS						75 DAS						105 DAS						At harvest					
	Kharif			Rabi			Kharif			Rabi			Kharif			Rabi			Kharif			Rabi		
MAIN PLOTS	2011	2012	Pooled	2011	2012	Pool	2011	2012	Pooled	2011	2012	Pooled	2011	2012	Pooled	2011	2012	Pooled	2011	2012	Pooled	2011	2012	Pooled
M1 (SRI)	1990	2607	2298	2290	3126	2708	4106	4647	4413	4490	5308	4899	6800	7736	7268	7081	8341	7711	9187	10161	9674	10082	11083	10582
M2 (Modified Drum seeder)	1556	2252	1904	1667	2547	2107	3521	3754	3676	3597	4645	4121	6383	7173	6778	6619	7558	7089	8400	9565	8983	9203	10381	9792
M3 ( Normal Drum seeder)	1761	1754	1758	2013	1861	1937	3232	3375	3284	3793	3967	3880	6265	6551	6408	6964	6683	6823	8104	8663	8383	9383	9477	9430
M4 (NTP)	1341	1481	1412	1669	1663	1667	2993	3050	3012	3325	3370	3348	6129	6053	6091	6148	6217	6182	7818	8223	8021	8783	8958	8870
S.E.M	66	48	50	52	61	45	86	86	71	64	109	51	130	163	114	148	169	52	212	259	189	190	267	136
C.D(P=.05)	229	167	174	180	214	157	297	297	247	223	378	177	451	564	395	512	583	181	732	895	653	656	922	471
SUBPLOTS																								
N1 (100% Inorg)	1706	2148	1927	2040	2460	2250	3664	3838	3681	4034.5	4498.8	4266.6	6580	7037	6808	7086	7381	7233	8719	9333	9026	9610	10191	9901
N2 (75% Inorg+25% org)	1540	1883	1711	1723	2073	1898	3302	3485	3404	3586	4103	3844	6311	6691	6501	6408	7022	6715	7991	8901	8446	8962	9713	9337
N3 (50% Inorg+50% org)	2113	2418	2265	2360	2871	2615	4007	4377	4203	4373	5037	4705	7033	7432	7232	7481	7913	7697	9285	9854	9570	10375	10695	10535
N4 (100%org)	1290	1647	1469	1516	1795	1655	2880	3125	3096	3213	3652	3432	5654	6354	6004	5837	6483	6160	7514	8524	8019	8503	9299	8901
S.E.M	35	65	37	77	61	51	94	97	52	154	111	105	185	192	127	162	190	106	176	213	145	339	232	188
C.D(P=.05)	103	191	107	225	177	149	274	283	151	449	325	306	541	561	371	473	555	311	514	621	422	990	676	550
MEAN	1662	2024	1843	1910	2300	2105	3463	3706	3596	3801	4322	4062	6394	6878	6636	6703	7200	6951	8377	9153	8765	9362	9975	9668
INTERACTIONS																								
Sub at same level of Main																								
S.E.M	70	131	73	154	122	102	188	194	103	307	223	210	371	384	254	324	380	213	352	425	289	678	463	377
CD (P=.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Main at same level of Sub																								
S.E.M	90	123	81	104	122	100	184	188	114	129	222	189	347	371	248	296	370	192	371	450	314	617	481	354
CD (P=.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS



**Table 4. Number of tillers m<sup>-2</sup> at various growth stages as influenced by establishment methods and nutrient management practices**

MAIN PLOTS	60 DAS						90 DAS					
	Kharif			Rabi			Kharif			Rabi		
	2011	2012	Pooled	2011	2012	Pool	2011	2012	Pooled	2011	2012	Pooled
M1 (SRI)	281.33	330.75	306.04	327.00	374.17	350.58	361.25	385.08	373.17	378.75	406.75	392.75
M2 (Modified Drum Seeder)	273.50	309.25	291.38	275.50	359.33	317.42	324.67	356.25	340.46	321.92	352.17	337.04
M3 ( Normal Drum seeder)	184.00	247.50	215.75	198.33	245.25	221.79	282.50	307.50	295.00	290.83	310.00	300.42
M4 (NTP)	145.50	236.83	191.17	180.00	218.42	199.21	270.75	297.00	283.88	283.58	297.17	290.38
S.E.M	6.99	6.83	6.14	9.06	11.51	8.62	7.23	9.68	7.20	7.52	10.66	6.95
C.D. (P=.05)	24.19	23.63	21.23	31.34	39.84	29.84	25.00	33.49	24.93	26.02	36.87	24.06
<b>SUBPLOTS</b>												
N1 (100% Inorg)	246.50	291.83	269.17	270.75	312.17	291.46	334.08	350.25	342.17	340.25	361.42	350.83
N2 (75% Inorg+25% org)	193.83	268.25	231.04	227.25	271.92	249.58	284.25	319.25	301.75	297.17	319.75	308.46
N3 (50% Inorg+50% org)	283.50	317.00	300.25	300.08	347.67	323.88	368.83	377.08	372.96	377.08	401.50	389.29
N4 (100%org)	160.50	247.25	203.88	182.75	265.42	224.08	252.00	299.25	275.63	260.58	283.42	272.00
S.E.M	8.77	8.96	5.75	6.21	10.81	5.83	9.42	11.48	8.35	5.19	11.33	6.38
C.D. (P=.05)	25.59	26.14	16.77	18.12	31.55	17.02	27.51	33.50	24.38	15.14	33.06	18.63
MEAN	221.08	281.08	251.08	245.21	299.29	272.25	309.79	336.46	323.13	318.77	341.52	330.15
<b>INTERACTIONS</b>												
<b>Sub at same level of Main</b>												
S.E.M	17.53	17.91	11.49	12.42	21.62	11.66	18.85	22.95	16.70	10.37	22.65	12.77
CD (P=.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
<b>Main at same level of Sub</b>												
S.E.M	16.72	16.95	11.69	14.06	21.98	13.28	17.85	22.11	16.16	11.72	22.33	13.06
CD (P=.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Table 5. Grain yield (kg ha<sup>-1</sup>) as influenced by establishment methods and nutrient management practices

Treatments	Kharif			Rabi		
	2011	2012	Pooled	2011	2012	Pooled
MAIN PLOTS	2011	2012	Pooled	2011	2012	Pooled
M1 (SRI)	6140	6535	6338	6438	6645	6541
M2 (Modified Drum seeder)	5356	6244	5800	5859	6287	6073
M3 ( Normal Drum seeder)	5084	5635	5359	5392	6090	5741
M4 (NTP)	4944	5468	5206	5356	6025	5690
S.E.M	137	138	34	112	117	47
C.D. ( <i>P</i> =.05)	475	477	117	388	405	162
SUBPLOTS						
N1 (100% Inorg)	5363	6124	5743	5898	6408	6153
N2 (75% Inorg+25% org)	5300	5827	5563	5591	6203	5897
N3 (50% Inorg+50% org)	5693	6383	6038	6033	6457	6245
N4 (100%org)	5168	5547	5358	5523	5980	5751
S.E.M	107	92	67	108	106	86
C.D. ( <i>P</i> =.05)	311	269	196	316	309	252
GENERAL MEAN	5381	5970	5676	5761	6262	6011
INTERACTIONS						
Sub at same level of Main						
S.E.M	213	185	134	216	211	173
CD ( <i>P</i> =.05)	622	539	391	631	617	505
Main at same level of Sub						
S.E.M	230	211	121	218	217	157
CD ( <i>P</i> =.05)	716	665	358	669	668	465

**Table 5a. Interaction of crop establishment methods and nutrient management practices on grain yield (kg ha<sup>-1</sup>)**

Establishment methods	Kharif 2011					Kharif 2012					Kharif pooled mean					Rabi 2011-12					Rabi 2012-13					Rabi pooled mean				
	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN
M1-SRI	6292	6092	6762	5414	6140	6974	6821	6523	5822	6535	6556	6307	6868	5618	6337	6534	6310	7090	5817	6438	7058	6646	7073	5801	6645	6796	6478	7081	5809	6541
M2- Modified drum seeder	5482	4942	5749	5249	5356	6643	6625	5872	583	6244	6053	5407	6196	5542	5799	6079	6018	6055	5283	5859	6363	6041	6603	6141	6287	6220	6029	6329	5711	6072
M3- Normal drum seeder	4753	5451	4910	5222	5084	6221	5929	5539	4851	5635	5341	5494	5565	5036	5359	5416	4953	5315	5882	5392	6433	5995	6023	5907	6090	5924	5474	5669	5894	5740
M4- NTP	4923	4714	5351	4787	4944	5695	5121	5375	5681	5468	5022	5044	5522	5233	5205	5562	5082	5670	5109	5356	5777	6128	6127	6069	6025	5669	5604	5898	5589	5690
<b>MEAN</b>	5363	5300	5693	5168		6383	6124	5827	5547		5743	5563	6038	5357		5898	5591	6033	5523		6408	6203	6457	5980		6152	5896	6244	5751	
	S.E.M		CD			S.E.M		(P=.05)			S.E.M		CD at			S.E.M		CD at			S.E.M		CD at			S.E.M		CD at		
Sub at same level of main	213		622			185		539			134		391			216		631			211		617			173		505		
Main at same level of sub	230		716			211		665			121		358			218		669			217		668			157		465		

**Table 6. Straw yield (kg ha<sup>-1</sup>) of rice as influenced by establishment methods and nutrient management practices means**

MAIN PLOTS	Kharif			Rabi		
	2011	2012	Pooled	2011	2012	Pooled
M1 (SRI)	7188.08	7542.75	7365.42	7342.08	7652.58	7497.33
M2 (Modified Drum Seeder)	6109.92	6956.08	6533.00	6679.17	7121.67	6900.42
M3 ( Normal Drum Seeder)	5793.08	6480.75	6136.92	6083.42	6539.33	6311.38
M4 (NTP)	5278.08	5974.58	5626.33	5759.00	6302.17	6030.58
S.Em±	140.83	185.79	123.59	155.42	195.07	126.46
C.D. at 5%	487.35	642.92	427.68	537.81	675.04	437.62
<b>SUBPLOTS</b>						
N1 (100% Inorg)	6138.33	6922.25	6530.29	6588.42	7044.25	6816.33
N2 (75% Inorg+25% org)	5996.17	6545.25	6270.71	6313.42	6749.17	6531.29
N3 (50% Inorg+50% org)	6617.25	7393.67	7005.46	7026.00	7486.33	7256.17
N4 (100%org)	5617.42	6093.00	5855.21	5935.83	6336.00	6135.92
S.Em±	131.83	158.55	92.02	147.15	170.44	106.65
C.D. at 5%	384.78	462.77	268.58	429.50	497.49	311.28
General Mean	6092.29	6738.54	6415.42	6465.92	6903.94	6684.93
<b>INTERACTIONS</b>						
<b>Sub at same level of Main</b>						
S.Em±	263.66	317.10	184.04	294.30	340.89	213.30
CD at 5%	NS	NS	NS	NS	NS	NS
<b>Main at same level of Sub</b>						
S.Em±	268.28	331.56	201.68	298.52	353.85	223.86
CD at 5%	NS	NS	NS	NS	NS	NS

**Table 7a. Interaction of crop establishment methods and nutrient management practices on LAI**

Establishment methods	at 45 DAS during kharif 2011					at 45 DAS in pooled kharif					at 75 DAS during kharif 2011					at 75 DAS during kharif 2012					at 75 DAS in pooled kharif				
	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN
M1-SRI	1.74	1.10	1.96	0.98	1.44	2.03	1.80	1.24	1.09	1.54	4.65	3.84	5.86	2.90	4.31	4.34	3.39	5.21	2.71	3.91	5.54	4.50	3.62	2.81	4.11
M2- Modified drum seeder	1.45	1.24	1.65	0.84	1.30	1.72	1.50	1.25	1.03	1.37	3.45	3.17	4.06	2.84	3.38	3.30	2.98	4.01	2.70	3.25	4.04	3.38	3.08	2.77	3.31
M3- Normal drum seeder	1.23	0.96	1.36	0.70	1.06	1.44	1.33	1.14	0.84	1.19	2.65	2.39	3.24	2.04	2.58	2.42	2.09	2.96	1.98	2.36	3.10	2.54	2.24	2.01	2.47
M4- NTP	0.90	0.71	1.10	0.65	0.84	1.25	1.10	0.88	0.75	1.00	2.36	1.96	2.74	1.78	2.21	2.36	2.07	2.67	1.84	2.24	2.71	2.36	2.02	1.81	2.22
<b>MEAN</b>	1.33	1.00	1.52	0.79		1.61	1.43	1.13	0.93		3.28	2.84	3.98	2.39		3.11	2.63	3.71	2.31		3.84	3.19	2.74	2.35	
	S.Em±	CD at 5%				S.Em±	CD at 5%				S.Em±	CD at 5%				S.Em±	CD at 5%				S.Em±	CD at 5%			
Sub at same level of main	0.06	0.16				0.04	0.12				0.16	0.47				0.16	0.48				0.13	0.38			
Main at same level of sub	0.06	0.17				0.04	0.12				0.16	0.47				0.16	0.49				0.13	0.39			

**Table 7b. Interaction of crop establishment methods and nutrient management practices on LA**

Establishment methods	75 DAS during rabi 2011-12					at 75 DAS in pooled rabi				
	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN	N <sub>1</sub>	N <sub>2</sub>	N <sub>3</sub>	N <sub>4</sub>	MEAN
M1-SRI	5.26	4.12	6.09	3.40	4.72	6.15	5.45	4.42	3.71	4.93
M2- Modified drum seeder	3.76	3.39	4.10	3.23	3.62	4.50	4.02	3.66	3.41	3.90
M3- Normal drum seeder	3.49	3.15	3.61	2.98	3.31	3.97	3.70	3.33	3.03	3.51
M4- NTP	2.86	2.70	3.09	2.12	2.69	3.26	3.00	2.78	2.35	2.85
<b>MEAN</b>	3.84	3.34	4.22	2.93		4.47	4.04	3.55	3.12	
	S.Em±		CD at 5%			S.Em±		CD at 5%		
Sub at same level of main	0.25		0.74			0.15		0.45		
Main at same level of sub	0.25		0.78			0.18		0.55		

#### 4. CONCLUSIONS

SRI registered superior growth characters and higher yield of rice over other establishment methods. Among nutrient management practices 50% inorganic+50% organic performed better in terms of growth and yield. A combination of SRI along with 50% inorganic+50% organic nutrient management practice recorded significantly superior growth characters and higher yield over the rest of the treatments.

#### ACKNOWLEDGEMENT

Authors express their gratefulness to Professor Jayashankar Telangana State Agricultural University and Indian Institute of Rice Research for providing funds and necessary facilities to conduct the research work.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Mohanty S, Yamano T. Rice food security in India: Emerging challenges and opportunities, The future rice strategy for India, Academic Press, Oxford. 2017;Chapter-1:1-13
- Baishya LK, Rathore SS, Dharmendrasingh D, Deka BC. Effect of integrated nutrient management on rice productivity, profitability and soil fertility. *Annals of Plant and Soil Research*. 2015;17(1):86–90.
- Chakraborty D, Ladha JK, Rana DS, Jat ML, Gathala MK, Yadav S, et al. A global analysis of alternative tillage and crop establishment practices for economically and environmentally efficient rice production. *Scientific reports*. 2017;7(1):1–11.
- Xu L, Li X, Wang X, Xiong D, Wang F. Comparing the grain yields of direct-seeded and transplanted rice: A meta-analysis. *Agronomy*. 2019;9(11):767.
- Guru P, Choudhary R, Kumar A, Jhorar R, Singh VD. Recent production technologies of rice for its sustainable cultivation in Haryana. *International Journal of Agricultural Sciences*. 2017;13(1): 83-92.
- Singh SK, Abraham T, Kumar R, Kumar R. Response of crop establishment methods and split application of nitrogen on productivity of rice under irrigated ecosystem. *Environment and ecology*. 2017;35(2A):859-862.
- Saravanane P. Effect of different weed management options on weed flora, rice grain yield and economics in dry direct-seeded rice. *Indian journal of weed science*. 2020;52(2):102-106.
- Bedari A, Rawat G, Amgain LP, Thapa DB, Poudel M. Productivity and profitability of different varieties of Chaiterice grown under different sowing methods at Baniyani, Jhapa. *Nepalese journal of agricultural sciences*. 2020; 19:155-164.
- Sahoo K, Ray M, Sahoo SK. Effect of different crop establishment methods on yield and yield of wet season rice. *Environment and ecology*. 2017; 35(3A):1854-1858.
- Rahman A, Salam MA, Kader MA. Effect of crop establishment methods on the yield of boro rice. *Journal of bangladesh agricultural University*. 2019;17(4):521–525.
- Kaur J, Mahal SS. Irrigation scheduling in direct seeded rice (*Oryza sativa* L.)-A review. *Agricultural Reviews*. 2015;36(3):208–217.
- Bhat MA, Hussain A, Ganai MA, Jehangir IA, Teli NA. Effect of different crop establishment methods and nutrient management practices on yield of rice under Kashmir valley conditions. *Journal of crop and weed*. 2018;14(1):30-34.
- Singh CS, Mohit SS, Singh K, Singh AK, Ashok KR. Growth and yield response of rice cultivars under system of rice intensification and conventional method of rice production system. *International journal for environmental research*. 2015; 9(3&4):1077–1081.
- Senthilvalavan P, Ravichandran M. Growth and physiological characters of rice as influenced by integrated nutrient management under SRI in Cauvery deltaic zone of Tamil Nadu. *Annals of plant and soil research*. 2019;21(3):210–216.
- Kumar V, Kumar T, Singh G, Singh RA. Effect of integrated nutrient management on yield of rice and its residual effect on

- wheat in rice-wheat system under lowland. *Annals of plant and soil research*. 2017;19(4):360-365.
16. Mondal S, Mallikarjun M, Ghosh M, Ghosh DC, Timsina J. Effect of integrated nutrient management on growth and productivity of hybrid rice. *Journal of Agricultural Science and Technology*. 2015;B5:297-308.
  17. Sarangi DR, Sahoo TR, Sethy S, Chourasia M, Prasad SM, Mohanta RK, et al. Effect of replacing a part of nitrogenous fertilizer by brown manuring in direct seeded rice: A field study. *Oryza- an international journal on rice*. 2016; 53(2):226-228.
  18. Aasif M, Chinnamani I, Kumar NS, Hemalatha M, Suresh, S. Influence of integrated nutrient management practices on yield and nutrient uptake of rice under system of rice intensification. *International journal of advances in agricultural science and technology*. 2018;5(7):10-16.
  19. Pandit TK, Mookherjee S, Karforma J. Performance of direct seeded rice under integrated nutrient management practices in old alluvial soils of West Bengal. *International research journal of pure & applied chemistry*. 2020;21(5):19-24.
  20. Singh D, Kumar A. Effect of sources of nitrogen on growth, yield and uptake of nutrient in rice. *Annals of plant and soil research*. 2014;16(4):359–361.
  21. Yadav L, Meena N. Performance of aromatic rice genotype as influenced by integrated nitrogen management. *Indian Journal of Agronomy*. 2014;59(2):251–255.
  22. Jat LK, Singh YV. Plant growth and yield as affected by application of organic inputs with fertilizer in rice wheat cropping sequence. *Indian journal of plant and soil*. 2019;6(1):25-31.
  23. Hussain A, Lone AH, Bhat MA, Ganai MA, Teeli NA, Najeeb S, Shikari AB. Yield and physiological response of newly released rice (*Oryza sativa*) varieties to crop establishment methods under temperate conditions of Kashmir. *Indian journal of agronomy*. 2018;63(4):441-446.
  24. Jat AL, Srivastava VK, Singh RK. Effect of crop establishment methods and integrated nitrogen management on productivity of hybrid rice-wheat cropping system. *Indian journal of agronomy*. 2015;60(3):341–346.
  25. Wahlang B, Das A, Layek J, Munda GC, Ramkrushna GI, Panwar AS. Effect of establishment methods and nutrient management on physiological attributes and water-use efficiency of rice in a sub-tropical climate. *Indian journal of agronomy*. 2015;60(4):534-540
  26. Thirumeninathan S, Jayanthi C, Karthikeyan R, Ravichandiran V. Effect of different crop establishment techniques on growth, yield attributes and yield of rice under puddled condition. *Madras Agricultural Journal*. 2017;104(10-12):315-318.
  27. Singh DK, Pandey PC, Ali N, Gupta S. Stand establishment techniques of rice in conjunction with nutrient sources for soil health and productivity of rice-wheat cropping system. *Indian journal of agronomy*. 2015b;60(1):31-37.
  28. Upendrarao A, Ramanamurthy KV, Madhukumar K, Visalakashmi V, Harisatyanarayana N, Govindarao S. Alternate crop establishment methods for water-saving and high rice productivity in north coastal Andhra Pradesh. *Current agriculture research journal*. 2020;8(3):219-223.
  29. Pramod K, Ghanshyam S, Prashantdeo S, Tejbal S, Anand S, Lakrakairovin. Effect of different crop establishment methods and nitrogen levels on the performance of kharif season rice in the Indo-Gangetic plains of eastern Uttar Pradesh. *Crop research*. 2021;56(1&2):1-7.
  30. Amanullah, Hidayatullah. Influence of organic and inorganic nitrogen on grain yield and yield components of hybrid rice in north western Pakistan. *Rice science*. 2016;23(6):326–333.
  31. MeherMalikaMd, Islam RMd, RezaulKarim, Huda A, Jahiruddin M. Organic and inorganic fertilizers influence the nutrient use efficiency and yield of a rice variety BINA dhan7. *Academic research journal of agricultural science and research*. 2015;3(7):192-200.
  32. Elhabe, H. Effect of organic and inorganic fertilizers on rice and some nutrients availability under different water regimes. *Journal of agricultural science and food research*. 2018;9(4):247-251.

33. Nahar L, Sarker ABS, Mahbub MM, Akter R. Effect of crop establishment method and nutrient management on yield and yield attributes of short duration aman rice. Bangladesh agronomy journal. 2017; 21(1):117-123.

---

© 2021 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/79936>