



Evaluation of Different Adenium (*Adenium arabicum*) Hybrids Performance for Seed Germination and Seedling Growth under Prayagraj Agro-climatic Conditions

Al-Fathima S^{a++*} and Urfi Fatmi^{a#}

^a Department of Horticulture, SHUATS, Prayagraj, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i74824>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119837>

Original Research Article

Received: 05/05/2024

Accepted: 08/07/2024

Published: 09/07/2024

ABSTRACT

An experiment was conducted in the Department of Horticulture, Sam Higginbottom University of Agriculture Technology And Sciences, Prayagraj, during the 2023-2024 period. The study employed Completely Randomized Design (CRD) with 10 hybrids, each replicated thrice. The hybrids tested were Godji x Nomsod, Godji x Ara Champ, Godji x Emerald Tank, Bangkla, Dynamo, Tropido, Dino King, Peth Dino, Susthaskorn, and Peth Phunlan. Among these hybrids the hybrid Godji x Ara Champ exhibited significantly better performance in several parameters, including germination percentage, days to 50% germination, seedling vigor index, seedling height, germination speed index, survival percentage, number of leaves per seedling, leaf area, caudex diameter. This hybrid's performance was found to be at par with the hybrid Dino King in terms of plant height, leaf area,

⁺⁺ M. Sc. Scholar;

[#] Assistant Professor;

^{*}Corresponding author: E-mail: alfathishalu@gmail.com;

Cite as: S, Al-Fathima, and Urfi Fatmi. 2024. "Evaluation of Different Adenium (*Adenium Arabicum*) Hybrids Performance for Seed Germination and Seedling Growth under Prayagraj Agro-Climatic Conditions". *International Journal of Plant & Soil Science* 36 (7):1092-97. <https://doi.org/10.9734/ijpss/2024/v36i74824>.

caudex diameter, survival percentage. Consequently, hybrids Godji x Ara Champ and Dino King demonstrated superior performance in both germination and seedling growth under the agro-climatic conditions of Prayagraj.

Keywords: *Adenium*; germination; growth; hybrids; seed.

1. INTRODUCTION

Adenium, a genus belonging to the family Apocynaceae, is renowned as a striking exotic houseplant, often recognized by its distinctive appearance resembling a bush or tree, commonly referred to as the desert rose. Highly favored for its low maintenance and ornamental appeal, *Adenium* holds significant relevance in the market due to its sculptural aspect, remarkable resistance to drought stress, glossy leaves, and strikingly vibrant flowers [1,2,3,4,5]. *Adenium arabicum*, a species of succulent plant, exhibits a striking resemblance to a miniature baobab tree, characterized by its grotesquely swollen trunk. While traditionally considered a distinct species, it is now recognized as a local or morphological form of *Adenium obesum*. Nonetheless, it remains highly popular among collectors of 'fat plants' [6].

Adenium arabicum shares similarities in appearance with *Adenium obesum*, but is distinguished by its notably massive, squat, and fat caudex, with minimal differentiation between trunk and branches. The bark of this species often displays hues ranging from purplish to dark brown, evoking a distinctively desert-like aesthetic. Its fleshy trunks and short, naked branches are further accentuated by a purple to dark brown coloration. Additionally, *Adenium arabicum* is characterized by larger, thick, fleshy leaves that boast a shiny surface on both the upper and lower sides. Despite variations in habit, most cultivated plants are hybrids, and only a few cultivar names have a formally recognized description.

Plants grown from seed often develop the most astonishing shapes of caudex within just 10 years of growth. Propagation by seeds is preferred to ensure the optimal development of the caudex, the distinctive swollen trunk characteristic of *Adenium* plants. Desert rose can be propagated via seeds or cuttings, with vegetative propagation methods including root cuttings, grafting, and air layering also being viable options. Modern hybrids of *Adenium* exhibit luxurious flowering, adding to their ornamental appeal. Even during dormancy, when the plant is not actively growing, its beauty is

retained due to the captivating shape of the caudex, often characterized by a thickening below the main trunk [7].

The propagation of the species occurs mainly by seeds, whereas plants from seeds present more developed caudex and main root when compared to the ones propagated by cuttings. To ensure optimal seed germination and seedling growth, selecting the appropriate substrate is crucial for these species when produced on a large scale. Although the easiest method of propagation is through cuttings, plants obtained via this method are often not well received in the ornamental market. This is because they tend to produce underground caudex and lack the same exuberance as plants propagated from seeds. Therefore, it is essential to prioritize seed propagation and evaluate the seedling growth of different *adenium* hybrids to ensure the propagation of species with well-developed caudex.

2. MATERIALS AND METHODS

The present investigation was carried out in the Department of Horticulture, Sam Higginbottom University of Agriculture Technology and Science, Prayagraj, during 2023-2024. The experimental material consisted of 10 *adenium* hybrids viz., Godji x Nomsod, Godji x Ara Champ, Godji x Emerald Tank, Bangkla, Dynamo, Tropido, Dino King, Peth Dino, Susthakorn and Peth Phunlan. whereas 15 number of plants per hybrids were used respectively. The experiment was laid out in Completely Randomized Design (CRD) with 10 hybrids and each hybrid was replicated thrice. The data recorded during the experiment were subjected to statistical analysis by using analysis of variance (ANOVA).

3. RESULTS AND DISCUSSION

3.1 Seed Germination Parameters

Among the seed germination parameters studied, significant variations were observed among the 10 *Adenium* hybrids and the data is presented in Table 1.

Table 1. Seed Germination parameters of different adenium hybrids

Hybrids	Germination percentage	Days to 50% germination (days)	Seedling vigour index (%)	Germination speed index
Godji x Nomsod	80	6.07	546.67	0.75
Godji x Ara Champ	100	5.33	1576.67	1.22
Godji x Emerald Tank	80	7.83	588.67	0.73
Bangkla	86	5.57	1043.33	0.96
Dynamo	73	6.27	606.67	0.86
Tropido	66	7.75	583.33	0.91
Dino King	93	5.80	1080.00	1.02
Peth Dino	60	8.40	440.00	0.77
Susthakorn	80	6.33	856.67	0.84
Peth phunlan	73	8.20	637.33	0.78
F- Test	S	S	S	S
SE(d)±	8.43	1.04	129.07	0.03
CD_(0.05)	17.59	2.17	269.23	0.07
CV(%)	13.02	18.82	19.86	4.63

Table 2. Vegetative parameters of different adenium hybrids

Hybrids	Seedling height(cm)	Number of leaves/ seedling	Leaf area (cm ²)	Caudex diameter (cm)	Survival Percentage (%)
Godji x Nomsod	9.9	18.6	10.73	2.83	73
Godji x Ara Champ	13.0	23.3	12.83	4.10	100
Godji x Emerald Tank	9.0	18.0	8.83	3.0	80
Bangkla	12.0	21.0	11.83	3.57	86
Dynamo	9.9	19.3	9.67	2.80	66
Tropido	9.2	19.0	8.20	3.17	66
Dino King	12.5	22.0	12.17	3.80	93
Peth Dino	7.6	16.6	5.37	1.17	53
Susthakorn	8.8	19.0	7.17	2.53	73
Peth phunlan	9.2	19.6	7.0	2.17	73
F- Test	S	S	S	S	S
SE(d)±	0.28	0.67	0.42	0.19	9.89
CD_(0.05)	0.59	1.39	0.88	0.40	20.63
CV(%)	3.42	4.15	5.54	8.05	15.80

Significantly higher germination percentage (100%) was observed in hybrid H₂ (Godji x Ara Champ) which was found to be at par with hybrid H₇ (Dino King, 93.33%) while lesser germination (60%) was observed in H₈ (Peth Dino). Variation in germination percentage could be influenced by the parental genotypes, their genetic makeup as well as prevailing temperature during the growing period. Similar results are recorded in adenium by [8]. Significantly, the lesser number of days to 50% germination (5.33) was observed in hybrid H₂ (Godji x Ara Champ) which was found to be at par with hybrid H₄ (Bangkla, 5.57) while more number of days to germination (8.40) was observed in H₈ (Peth Dino). The variation in germination days can be influenced by the genetic makeup of the parental genotypes and

the environmental conditions during the growing period. Similar results have been recorded in adenium by [8] indicating that these factors play a significant role in germination percentages. Similar findings were observed in [9,10,11,12] and [13].

Significantly, higher seedling vigour index (1576) was observed in hybrid H₂ (Godji x Ara Champ) which was found to be at par with hybrid H₇ (Dino King, 1080) while lesser seedling vigour index (440) was observed in H₈ (Peth Dino) Variation in seedling vigour index depends upon the shoot length of the seedlings and influenced by the parental genotypes, their genetic makeup as well as prevailing temperature during the growth period. Similar results recorded in tuberose by

[14] Significantly, higher germination speed index (1.22) was observed in hybrid H₂ (Godji x Ara Champ) which was found to be at par with hybrid H₇ (Dino King, 1.02) while lesser germination speed index (0.77) was observed in H₈ (Peth Dino) Difference in germination speed index may be due to the inheritant character and genetic makeup of the hybrids and the prevailing environmental conditions. Similar results were recorded in adenium by [15].

3.2 Vegetative Parameters

From the present investigation, it is concluded that significant variations were observed among the 10 Adenium hybrids studied across all the vegetative parameters and the data is presented in Table 2.

Significantly, higher seedling height (13.03) was observed in hybrid H₂ (Godji x Ara Champ) which was found to be at par with hybrid H₇ (Dino King, 12.57) while lesser seedling height (7.67) was observed in H₈ (Peth Dino) Difference in seedling height can be attributed to their genetic makeup and environmental conditions prevailed. Similar results were recorded in desert rose by [16]. Significantly, more number of leaves (23.33) was observed in hybrid H₂ (Godji x Ara Champ) which was found to be at par with hybrid H₇ (Dino King, 22) while lesser number of leaves (16.67) was observed in H₈ (Peth Dino). The variation in number of leaves per plant among the hybrids might be due to the rate of vegetative growth among the hybrids that could be attributed to genetic makeup and could have been further influence by environmental conditions. Similar results were recorded in adenium by [17].

Significantly maximum leaf area (12.83) was observed in hybrid H₂ (Godji x Ara Champ) which was found to be at par with hybrid H₇ (Dino King, 12.17) while minimum leaf area (5.37) was observed in H₈ (Peth Dino) leaf area of all the hybrids were recorded and significant differences were observed at 360 days after transplanting. Significant variation in leaf area can be attributed to parental genotypes of the hybrids along with environmental conditions. Similar results were shown in adenium by [7]. Significantly, higher survival percentage (100%) was observed in hybrid H₂ (Godji x Ara Champ) which was found to be at par with hybrid H₇ (Dino King, 93%) while lesser survival percentage (53%) was observed in H₈ (Peth Dino). Germinated seeds are highly vulnerable to lack of moisture for

growth, fire, herbivores, burial under litter, being washed away by rain, and heat on bare soil, and hence up to 90 % of released seed will not make it past the seedling stage were recorded in adenium by [6]. Significantly maximum caudex diameter (4.10) was observed in hybrid H₂ (Godji x Ara Champ) which was found to be at par with hybrid H₇ (Dino King, 3.80) while lesser germination (1.17) was observed in H₈ (Peth Dino) caudex diameter of all hybrids were recorded and significant differences were observed at 360 days after transplanting. Significant variation in caudex diameter can be attributed to genetic variability of the hybrids along with environmental conditions which govern the plant growth were recorded in adenium by [18] and [19-22].

4. CONCLUSION

From the present investigation, it is concluded that the 10 Adenium hybrids under study exhibited significant variation across all observed parameters. The hybrid Godji x Ara Champ demonstrated superior performance in parameters such as germination percentage, days to 50% germination, seedling vigour index, seedling height, germination speed index, survival percentage, number of leaves per seedling, leaf area, caudex diameter. Additionally, its performance was comparable to that of the hybrid Dino King in plant height, leaf area, caudex diameter, survival percentage. Therefore, the hybrids Godji x Ara Champ and Dino King showed exceptional performance in both germination and seedling growth under the agro-climatic conditions of Prayagraj.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Dimmitt M, Edwards T. Adenium taxonomy and nomenclature: Progress creates more questions. *Cactus and Succulent Journal*. 2021;93(4):252-266.

2. Wani, Aadil Akbar, Faisal Ur Rasool, Latief Ahmad, Mohammad Iqbal Bhat, Kahkashan Qayoom, Aamir Hassan Mir, Raies A Bhat. Effect of planting density and nitrogen levels on the yield attributing characters of fodder maize (Shalimar Fodder Maize-1)". Asian Journal of Soil Science and Plant Nutrition. 2024;10(2):483-88. Available:<https://doi.org/10.9734/ajsspn/2024/v10i2306>
3. Tano, Ekra Kouamé, Oulo Alla N'nan, Jean-Noël Konan, Désiré Allou, Eugène Kouakou Konan, Assanvo Simon-Pierre N'guetta. Evaluation of the morphological quality of hybrid seeds and the vegetative development of plants from the two reproductive systems of elite dura genitors from oil palm (*Elaeisguineensis Jacq*) seed fields in Côte d'Ivoire". Journal of Experimental Agriculture International. 2021;43(10):64-72. Available:<https://doi.org/10.9734/jeai/2021/v43i1030748>
4. Hu Q, Fu Y, Guan Y, Lin C, Cao D, Hu W, Sheteiwy M, Hu J. Inhibitory effect of chemical combinations on seed germination and pre-harvest sprouting in hybrid rice. Plant Growth Regulation. 2016 Dec;80:281-9. Available:<https://link.springer.com/article/10.1007/s10725-016-0165-z>
5. Landbo L, Jørgensen RB. Seed germination in weedy Brassica campestris and its hybrids with B. napus: implications for risk assessment of transgenic oilseed rape. Euphytica. 1997 Oct;97:209-16. Available:<https://link.springer.com/article/10.1023/A:1003032804340>
6. Van der Walt K, Witkowski ETF. Seed viability, germination and seedling emergence of the critically endangered stem succulent, *Adenium swazicum*, in South Africa. South African Journal of Botany. 2017;109:237-245.
7. Paul D, Biswas K, Sinha SN. Biological activities of *Adenium obesum* (Forssk.) Roem. & Schult: a concise review. Malaya Journal of Biosciences. 2015;2(4):214.
8. Baskin JM, Baskin CC. A classification system for seed dormancy seed science research. 2004;14:1-16.
9. Snow AA, Moran-Palma P, Rieseberg LH, Wszelaki A, Seiler GJ. Fecundity, phenology, and seed dormancy of F₁ wild-crop hybrids in sunflower (*Helianthus annuus*). American Journal of Botany. 1998;85(6):794-801.
10. Archana B, Paul AA, Hunje R, Patil VS. Studies on genetic variability analysis in gladiolus hybrids. Journal of ornamental Horticulture. 2008;11(2):121-126.
11. Kumar PH, Kulkarni BS. Genetic variability in gladiolus for growth and flowering characters (*Gladiolus hybridus*). Journal of Horticultural Sciences. 2009;177-180.
12. Weiss AN, Primer SB, Pace BA, Mercer KL. Maternal effects and embryo genetics: Germination and dormancy of crop-wild sunflower hybrids. Seed Science Research. 2013;23(4):241-255.
13. Naresh S, Rao AD, Baskhar VV, Rao MP, Krishna KU. Genetic variability, heritability and genetic advance in gladiolus hybrids Plant Archives. 2015;15(1):377-381.
14. Ranchana P, Kannan M. Self and cross compatibility studies in tuberose (*Pollianthes tuberosa*). Journal of Genetics and Plant Breeding. 2016;11(1):33-36.220.
15. Colombo RC, Favetta V, Yamamoto LY, Alves GAC, Abati J, Takahashi LSA, De Faria RT. Biometric description of fruits and seeds, germination and imbibition pattern of desert rose [*Adenium obesum* (Forssk.), Roem. & Schult.] Journal of Seed Science. 2015;37(4):206-213.
16. Colombo RC, Favetta V, De Carvalho DU, Da Cruz MA, Roberto SR, De Faria RT. Production of desert rose seedlings in different potting media. Ornamental Horticulture. 2017;23(3):250-256.
17. Dimmit M, Hanson C. The genus adenium in cultivation (*A. obesum* and multiflorum). Cactus and Succulent Journal. 2009;63(5):223-225.
18. Dimmitt MG. Adenium culture, growing large specimens quickly. Cactus Succulent Journal. 1998;63(5):59-64.
19. Varella TL, Silva GM, Maximiliano KZ, Mikovski AI, Silva Carvalho IF, Silva ML. *In vitro* germination of desert rose varieties Ornamental Horticulture. 2015;21(2):227-234.
20. Cabral JM, Melo LD, Junior JL, Berto T, Crisostomo NM, Gomes LC, Silva KW. Seed biometry, growth, and seedling development of *Adenium obesum* (Forssk.) Roem. & Schult. in substrates. GeSec: Revista de Gestaoe Secretariado. 2024;15(3):3569-3572.

21. Fleck IM, Rosset JS, Fleck L, Zoz T, Ozório JMB. Production of *Adenium obesum* (Forssk.) Roem. & Schult seedlings in different substrates. *Revista De Agricultura Neotropical*. 2024;11(2):8300-8304.
22. Kim SY, Lee HY, Park C, Kim D, Kim JB, Kim SH, Kim TS. Assessing amounts of genetic variability in key horticultural traits underlying core korean breeding lines of cut chrysanthemums. *Plants*. 2024;13(5):577-581.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119837>