



## **Correlation and Path Coefficient Analysis of Some Rice Genotypes (*Oryza sativa* L.)**

**Shraddha Singh<sup>a\*</sup>, A. K. Singh<sup>a</sup>, Sumant Pratap Singh<sup>b</sup>, Alok Kumar Singh<sup>a</sup> and R. K. Yadav<sup>a</sup>**

<sup>a</sup> Department of Crop Physiology, Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya - 224229 (U.P.), India.

<sup>b</sup> Department of Plant Molecular Biology and Genetic Engineering, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya - 224229 (U.P.), India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJECC/2022/v12i1030905

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/88495>

**Original Research Article**

**Received 10 April 2022**

**Accepted 20 June 2022**

**Published 21 June 2022**

### **ABSTRACT**

An experiment was carried out to study the correlation and path analysis in twenty-six rice (*Oryza sativa* L.) varieties, among 26 rice genotypes, 15 rice varieties and 11 landraces collections from hilly region of Mirzapur in eastern Uttar Pradesh. The experiment was performed during *kharif* 2021-22 in Randomized Block Design with three replications to analyze correlation and path Analysis. At phenotypic and genotypic level, number of grain per panicle (GPP 0.603), chlorophyll content (Chl 0.326), Ear bearing tillers per plant (EBT 0.356) showed highly positive significant correlation with yield per plant (YPP) to emerge as most important associates of grain yield in rice. Path analysis identified the highest positive direct effect on grain yield per plant was exhibited by number of grain per panicle (GPP 0.485), days to 50% flowering (DFF 0.414), plant height (PH at 45DAT 0.255), chlorophyll content (Chl 0.255), Ear bearing tillers per plant (EBT 0.173), plant height (PH at 30DAS 0.042), number of tillers per plant (NOT 0.029), panicle length (PL 0.024) are the most significant direct as well as indirect effect. Yield contributing components which under consideration at time of devising selection strategy aimed at developing varieties having higher yield.

**Keywords:** Correlation; path analysis; significant; phenotypic; genotypic landraces; rice.

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) occupies a pivotal place in Indian agriculture, as it forms the staple food for two-thirds of the population and provides 43 per cent calories requirement and 20-25% agriculture income. More than 90 percent of the world's rice is grown and consumed in Asia, where 60 per cent of the earth's people and two third of world's poor live [1]. Rice farming is about 10000-year-old and largest single use of land for producing food. About 11% of total Earth's arable land was covered by rice fields. The frequent occurrence of drought as well as other abiotic stresses has been identified as the major issue to the low productivity of rice in rainfed ecosystems, particularly in eastern region of India. Most of agronomical traits are quantitative traits showing normal distributions in phenotype of the traits. Information on association of characters, direct and indirect effects contributed by each character towards yield will be an added advantage in helping the selection process. Correlation and path analysis establish the extent of association between yield and yield components and also bring out relative importance of their direct and indirect effects, thus giving an obvious understanding of their association with grain yield [2]. Ultimately, this kind of analysis could help the breeder to design his selection strategies to improve grain yield. In the light of the above scenario, the present investigation is carried out with the objective of studying the character associations in rice for yield improvement.

## 2. MATERIALS AND METHODS

The present experiment was carried out at Student's Instructional Farm (SIF), Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya, India. Seeds of the 26 genotypes were sown in raised nursery bed. The seedlings were transplanted to the main field at the rate of one seedling per hill, after 21 days, with a spacing of 20cm x 15cm. The experiment was arranged in a randomized block design (RBD) with thrice replications. The recommended agronomical practices and plant protection measures were followed to ensure a normal crop growth and development. Observations were recorded on five randomly selected plants in each replication from the two centre rows. 14 traits viz. Days to flowering Initiation (DFIT), days to 50% flowering (DFF), days to 100% flowering (DHF), Plant height at the time of 15, 30, 45, Maturity (PH), no. of tillers (NOT), panicle length (PL), Spiklete's per

panicle (SPP), ear bearing tillers (EBT), grain per panicle (GPP), Chlorophyll (Chl), Grain yield per plant (GYP). Observations were recorded and the data was subjected to statistical analysis. Statistical analyses for the above characters were done following Singh and Chaudhary [3] for correlation coefficient and Dewey and Lu [4] for path analysis.

## 3. RESULTS AND DISCUSSION

In the present investigation for phenotypic correlation coefficients, the genotypic correlation coefficients were generally higher than their respective phenotypic correlation coefficients (Table 2). At phenotypic level, YPP showed highly positive significant correlation by GPP (0.603), Chl (0.326). Similar trends of results were also reported by Kishore et al. 2007. Positive significant correlation was obtained in EBT (0.249). Highly negative significant correlation was showed with PH at the time of maturity (0.341). Negative significant correlation was showed by PH at the time of 45DAT (0.223). Positive non-significant correlation was showed with DFF (0.149), DHF (0.102), DTFI (0.095), PL (0.088), SPP (0.072), PH at the time of 30DAS (0.006). GYP showed negative non-significant correlation with PH at the time of 15DAS (0.012), NOT (0.088). Chl content showed highly negative significant correlation by PH at the time of 15DAS, Positive significant correlation with GPP. Positive non-significant correlation with PL, SPP and PH at the time of maturity, PH at the time of 30 DAS, PH at the time of 45 DAT showed with negative non-significant correlation. GPP showed highly negative significant correlation with PH at the time of maturity, NOT. Positive non-significant correlation was obtained in GPP, PH at the time of 15DAS, PH at the time of 45DAT. Negative non-significant correlation in DHF, followed by EBT, PH at the time of 30DAS and EBT showed positive significant correlation in DHF. Positive non-significant correlation with DFF, PH at the time of 15DAS, DTFI, SPP, PH at the time of 45DAT, PH at the time of 30DAS, PH at the time of maturity, PL. SPP showed highly negative significant correlation in PH at the time of maturity and Negative significant correlation was showed with PH at the time of 45DAT. Positive non-significant correlation was obtained in NOT, DHF, DFF, DTFI, PL. Negative non-significant was obtained in PH at the time of 30DAS, PH at the time of 15DAS. PL showed positive significant correlation in PH at the time of 45DAS. Positive non-significant correlation was obtained in NOT, DFF, PH at the time of 30DAS,

DTFI, PH at the time of maturity. Negative non-significant correlation was obtained in PH at the time of 15DAS, DHF. Similar results were reported by Lalitha and Shreedhar (1996). NOT showed positive non-significant correlation in PH at the time of maturity. Negative non-significant correlation was obtained in PH at the time of 30DAS, PH at the time of 45DAT, DTFI, DHF, DFF, PH at the time of 15DAS. PH at the time of maturity showed highly positive significant correlation in DTFI, PH at the time of 45DAT, DFF. Positive significant correlation was obtained in PH at the time of 15DAS, DHF. Positive non-significant correlation was obtained in PH at the time of 30DAS. PH at the time of 45DAT showed highly positive correlation in DHF. Positive significant correlation was showed in PH at the time of 15DAS. Positive non-significant correlation was obtained in DTFI, DFF. PH at the time of 30DAS showed highly positive significant correlation in DHF. Positive significant correlation was obtained in PH at the time of 15DAS. Positive non-significant correlation was obtained in DTFI, DFF(0.054). PH at the time of 15DAS showed highly positive significant correlation with DFF, DHF. DTFI showed positive significant correlation. DHF showed highly positive significant correlation with DTFI, DFF. DFF showed highly positive significant correlation with DTFI. The positive association between these characters have also been reported by Chand *et al.*, 2007, Borbora *et al.* [5], Dhurai *et al.* [6], Namita *et al.* [7], Nikhil *et al.* [8] and Sathisha *et al.* [9].

### 3.1 Genotypic Correlation Coefficient

Yield per plant showed highly positive significant correlation with GPP (0.637), Chl (0.417), EBT (0.356). Highly negative significant correlation was obtained in PH at the time of maturity (0.375). Negative non-significant correlation was obtained with PH at the time of 45DAT (0.251), NOT (0.246). Positive non-significant correlation was showed by PL (0.199), DFF (0.154), DHF (0.122), SPP (0.117), DTFI (0.100). Negative non-significant correlation was obtained in PH at the time of 15DAS (0.002), PH at the time of 30DAS (0.070). Chl showed highly positive significant correlation by GPP (0.280). Positive significant correlation with PH at the time of 15DAS, PH at the time of 30DAS, DHF, DTFI (0.339), DFF, NOT and positive non-significant correlation was showed by EBT, PL, SPP. Negative non-significant correlation was obtained with PH at the time of maturity, PH at the time of 45DAT. Characters mentioned above has also

being reported in rice by earlier workers [10,11] GPP showed highly positive significant correlation was obtained in EBT and negative highly significant correlation was obtained in NOT, PH at the time of 45DAT, PH at the time of maturity. Positive non-significant correlation was obtained in PH at the time of 30DAS, SPP, DHF. Negative non-significant correlation was found in DFF, DTFI, PH at the time of 15DAS, PL. EBT showed highly positive significant correlation in PH at the time of 30DAS, PH at the time of 15DAS, DHF, DTFI, DFF, SPP, PH at the time of 45DAT. Negative non-significant correlation was obtained in NOT, PL. Positive non-significant correlation was obtained in PH at the time of maturity (0.082). SPP showed highly positive significant correlation in NOT. Highly negative significant correlation was found in PH at the time of 30DAS, PH at the time of maturity, PH at the time of 45DAT. Positive non-significant correlation was obtained in PL, DFF, DTFI, DHF. Negative significant correlation was obtained in PH at the time of 15DAS and PL showed highly significant correlation was obtained in PH at the time of 45DAT. Highly negative significant correlation was obtained in PH at the time of 30DAS. Negative significant correlation was obtained in NOT. Positive non-significant correlation was obtained in DFF, DTFI, PH at the time of maturity. Negative non-significant correlation was obtained in PH at the time of 15DAS, DHF. NOT showed highly negative significant correlation was obtained in PH at the time of 30DAS(1.088), PH at the time of 15DAS, DFF, DHF. Positive non-significant correlation was obtained with PH at the time of maturity. Negative non-significant correlation was found in PH at the time of 45DAT, DTFI. PH at the time of maturity showed highly positive significant correlation in PH at the time of 45DAT, PH at the time of 15DAS, DTFI, DFF. Positive significant correlation was showed in DHF, PH at the time of 30DAS. PH at the time of 45DAT showed highly positive significant correlation was obtained in PH at the time of 30DAS, PH at the time of 15DAS. Positive significant correlation was obtained in DFF. Positive non-significant correlation was obtained in DHF, DTFI. PH at the time of 30DAS showed highly positive significant correlation in DHF, PH at the time of 15DAS. Positive significant correlation was obtained in DTFI, DFF. In PH at the time of 15DAS showed highly positive significant correlation with DTFI, DFF, DHF. DHF showed highly positive significant correlation with DTFI, DFF. DFF showed highly positive significant correlation with DTFI [12], Chand *et al.* 2007.

**Table 1. Detailed list of selected rice genotypes and their origin**

<b>Name of variety</b>	<b>Parentage</b>	<b>Year of release</b>	<b>Duration (in days)</b>	<b>Eco-System</b>	<b>Salient Features</b>	<b>Recommended for cultivation</b>
Pusa Basmati-1	Pusa-150 x Karnal Local	1989	135	Irrigated Areas	Semi dwarf (85-95 cm), grains: super fine aromatic, Yield: 45 Q/ha.	Goa, Mizoram and Uttar Pradesh
NDR-359	BG-90-2-4 x 08677	1994	115-125	Irrigated Areas	Semi dwarf (90-95 cm), grains: short tipped, Yield: 50 Q/ha.	Uttar Pradesh, Bihar and Orissa
Pusa 1121 (Pusa Sugandha-5)		2005	135-140	Irrigated saline soils	Medium (97.3 cm); Grain-slender, mod. resit. to RTV, sheath rot, & BLB; Yield: 55-65 q/ha.	AP and Kerala
DRR Dhan-44	IR93376-B-B-130	2014	120	Upland and drought prone	Drought tolerant, HYV, Semi tall	Uttrakhand, Haryana and Bihar
Sahbhagi dhan		2009	105	Rainfed upland/lowland	LB grain, tolerant to drought. Res to leaf blast, mod. res to brown spot, sheath rot, SB and leaf folder, 3.5-4 t/ha	Orissa and Jharkhand
Swarna Sub-1	Swarna 3/IR 49830-7-1-2-3	2009	145	Flood prone shallow lowlands	Plant height- 83.3cm Tolerant to complete submergence between 15-17 days, 5.2 t/ha yield	UP, Uttrakhand, Haryana and Bihar
NDR 2064	Pant Dhan 4/Saket4 // NDR 2017	2007	115	irrigated areas	50-55 Q/ha yield grains are (M.S)Medium Size, High yielding	UP, Orissa and West Bengal
NDR 2065	Pant Dhan 4/Saket 4// NDR 2018	2011	120-125	irrigated areas	50-55Q/ha grains are LB(Long Bold), High yielding	UP, Orissa and West Bengal
NDR 97	Nanina-22 x Ratna	1992	90-95	Rain fed Uplands	Dwarf (75-80 cm), Yield: 25-30 Q/ha.	UP, Orissa and West Bengal
IR-64	IR-5857-33-2-1 x IR-2061-465-1-5-5	1991	115-120	Irrigated Areas	Semi dwarf (100 cm), grains: Yield: 58 Q/ha.	All India
Sarjoo-52	T(N)1 x Kashi	1982	130-133	Irrigated	Semi dwarf (98 cm), erect, grains: long bold, white, moderately resistant to Bacterial Leaf Blight, Yield: 50-60 Q/ha.	Uttar Pradesh.

Table 2. Estimation of phenotypic correlation coefficient for 14 characters in selected rice germplasm

Traits	DTFI	DFF	DHF	PH at the time of 15 DAS	PH at the time of 30 DAS	PH at the time of 45 DAT	PH at the time of Maturity	NOT	PL	SPP	EBT	GPP	Chl	GYP
DTFI	<b>1.000</b>	0.968**	0.865**	0.268*	0.068	0.187	0.315**	-0.041	0.051	0.079	0.172	-0.039	-0.307**	0.095
DFF		<b>1.000</b>	0.832**	0.322**	0.054	0.224*	0.305**	-0.090	0.136	0.112	0.184	-0.010	-0.275*	0.149
DHF			<b>1.000</b>	0.300**	0.311**	0.206	0.255*	-0.072	-0.034	0.114	0.246*	0.012	-0.301**	0.102
PH at the time of 15 DAS				<b>1.000</b>	0.262*	0.484**	0.288*	-0.171	-0.018	-0.188	0.177	-0.033	-0.342**	-0.012
PH at the time of 30 DAS					<b>1.000</b>	0.222*	0.107	-0.037	0.057	-0.071	0.061	0.012	-0.169	0.006
PH at the time of 45 DAS						<b>1.000</b>	0.740**	-0.039	0.237*	-0.283*	0.086	-0.480**	-0.189	-0.223*
PH at the time of Maturity							<b>1.000</b>	0.002	0.024	-0.347**	0.022	-0.439**	-0.160	-0.341**
NOT								<b>1.000</b>	0.201	0.193	-0.124	-0.120	-0.009	-0.088
PL									<b>1.000</b>	0.046	0.001	-0.133	0.051	0.088
SPP										<b>1.000</b>	0.170	-0.023	0.045	0.072
EBT											<b>1.000</b>	0.117	-0.051	0.249*
GPP												<b>1.000</b>	0.251*	0.603**
Chl													<b>1.000</b>	0.326**

Table 3. Estimation of genotypic correlation coefficient for 14 characters in selected rice germplasm

Traits	DTFI	DFF	DHF	PH at the time of 15 DAS	PH at the time of 30 DAS	PH at the time of 45 DAT	PH at the time of Maturity	NOT	PL	SPP	EBT	GPP	Chl	GYP
DTFI	<b>1.000</b>	0.972**	0.875**	0.307**	0.171	0.188	0.319**	-0.185	0.062	0.123	1.130**	-0.043	-0.339**	0.100
DFF		<b>1.000</b>	0.845**	0.379**	0.088	0.230*	0.309**	-0.367**	0.202	0.127	1.121**	-0.009	-0.306**	0.154
DHF			<b>1.000</b>	0.353**	0.755**	0.205	0.262*	-0.300**	-0.014	0.122	1.752**	0.010	-0.340**	0.122
PH at the time of 15 DAS				<b>1.000</b>	0.481**	0.561**	0.343**	-0.455**	-0.004	-0.166	1.986**	-0.080	-0.485**	-0.002
PH at the time of 30 DAS					<b>1.000</b>	0.592**	0.229*	-1.088**	-0.423**	-0.589**	2.599**	0.018	-0.390**	-0.070
PH at the time of 45 DAS						<b>1.000</b>	0.757**	-0.173	0.326**	-0.374**	0.561**	-0.509**	-0.203	-0.251*
PH at the time of Maturity							<b>1.000</b>	0.123	0.036	-0.485**	0.082	-0.457**	-0.183	-0.375**
NOT								<b>1.000</b>	-0.284*	0.978**	-1.707**	-0.570**	-0.280**	-0.246*
PL									<b>1.000</b>	0.204	-0.429**	-0.169	0.138	0.199
SPP										<b>1.000</b>	0.592**	0.014	0.042	0.117
EBT											<b>1.000</b>	1.121**	0.177	0.356**
GPP												<b>1.000</b>	0.280**	0.637**
Chl													<b>1.000</b>	0.417**

Table 4. Phenotypic Path Coefficient for 14 character in selected rice germplasm

Trait	DTFI	DHF	DHF	PH at the time of 15 DAS	PH at the time of 30 DAS	PH at the time of 45 DAS	PH at the time of maturity	NOT	PL	SPP	EBT	GPP	Chl	GYP
DTFI	<b>-0.055</b>	0.343	-0.095	-0.015	0.003	0.047	-0.116	-0.001	0.001	-0.005	0.030	-0.019	-0.078	0.095
DFF	-0.053	<b>0.414</b>	-0.092	-0.019	0.002	0.057	-0.112	-0.003	0.003	-0.007	0.032	-0.005	-0.070	0.149
DHF	-0.048	0.343	<b>-0.110</b>	-0.017	0.013	0.052	-0.094	-0.002	-0.001	-0.007	0.042	0.006	-0.077	0.102
PH at the time of 15 DAS	-0.015	0.133	-0.034	<b>-0.058</b>	0.011	0.122	-0.106	-0.005	0.000	0.011	0.031	-0.016	-0.087	-0.012
PH at the time of 30 DAS	-0.004	0.022	-0.034	-0.015	<b>0.042</b>	0.056	-0.039	-0.001	0.001	0.004	0.010	0.006	-0.043	0.006
PH at the time of 45 DAT	-0.010	0.088	-0.024	-0.028	0.009	<b>0.252</b>	-0.272	-0.001	0.006	0.017	0.015	-0.233	-0.048	-0.223*
PH at the time of maturity	-0.012	0.129	-0.031	-0.017	0.005	0.187	<b>-0.367</b>	0.000	0.001	0.020	0.004	-0.213	-0.041	-0.341**
NOT	0.002	-0.037	0.008	0.010	-0.002	-0.010	-0.001	<b>0.029</b>	0.005	-0.011	-0.021	-0.058	-0.002	-0.088
PL	-0.003	0.056	0.005	0.001	0.002	0.060	-0.009	0.006	<b>0.024</b>	-0.003	0.000	-0.064	0.013	0.088
SPP	-0.005	0.043	-0.013	0.011	-0.003	-0.071	0.127	0.006	0.001	<b>-0.058</b>	0.029	-0.011	0.012	0.072
EBT	-0.009	0.087	-0.027	-0.010	0.003	0.022	-0.008	-0.004	0.000	-0.010	<b>0.173</b>	0.057	-0.013	0.249*
GPP	0.002	-0.003	-0.001	0.002	0.001	-0.121	0.161	-0.004	-0.003	0.001	0.020	<b>0.485</b>	0.064	0.603**
Chl	0.017	-0.115	0.033	0.020	-0.007	-0.048	0.059	0.000	0.001	-0.003	-0.009	0.122	<b>0.255</b>	0.326**

*R square = 0.5292; Residual effect = 0.6861*

Table 5. Genotypic path coefficient for 14 character in selected rice genotypes

Traits	DTFI	DFF	DHF	PH at the time of 15 DAS	PH at the time of 30 DAS	PH at the time of 45 DAS	PH at the time of maturity	NOT	PL	SPP	EBT	GPP	Chl	GYP
DTFI	<b>1.034</b>	-0.598	-0.576	-0.051	0.017	-0.032	0.060	0.057	0.018	0.044	0.152	-0.016	-0.016	0.100
DFF	1.004	<b>-0.619</b>	-0.546	-0.063	0.009	-0.039	0.058	0.112	0.060	0.046	0.151	-0.004	-0.014	0.154
DHF	0.905	-0.525	<b>-0.648</b>	-0.059	0.076	-0.035	0.049	0.092	-0.004	0.044	0.236	0.004	-0.016	0.122
PHat the time of 15 DAS	0.308	-0.237	-0.233	<b>-0.167</b>	0.048	-0.095	0.064	0.139	-0.001	-0.059	0.268	-0.030	-0.023	-0.002
PHat the time of 30 DAS	0.176	-0.056	-0.492	-0.080	<b>0.100</b>	-0.101	0.043	0.332	-0.125	-0.211	0.350	0.007	-0.018	-0.070
PHat the time of 45 DAS	0.188	-0.145	-0.143	-0.094	0.059	<b>-0.170</b>	0.142	0.053	0.096	-0.134	0.076	-0.189	-0.009	-0.251*
PHat the time of Maturity	0.330	-0.191	-0.170	-0.057	0.023	-0.129	<b>0.187</b>	-0.038	0.011	-0.174	0.011	-0.170	-0.009	-0.375**
NOT	-0.192	0.243	0.195	0.076	-0.109	0.029	0.023	<b>-0.306</b>	-0.084	0.350	-0.230	-0.212	-0.013	-0.246*
PL	0.064	-0.125	0.009	0.001	-0.042	-0.055	0.007	0.087	<b>0.295</b>	0.073	-0.058	-0.063	0.006	0.199
SPP	0.132	-0.083	-0.079	0.028	-0.059	0.063	-0.091	-0.299	0.060	<b>0.358</b>	0.080	0.005	0.002	0.117
EBT	1.173	-0.644	-1.143	-0.332	0.261	-0.095	0.015	0.522	-0.127	0.212	<b>0.135</b>	0.417	0.008	0.356**
GPP	-0.045	0.006	-0.067	0.013	0.002	0.087	-0.086	0.174	-0.050	0.005	0.151	<b>0.372</b>	0.013	0.637**
Chl	-0.364	0.193	0.220	0.081	-0.039	0.034	-0.034	0.086	0.041	0.015	0.024	0.104	<b>0.046</b>	0.417**

R square = 0.3750; Residual effect = 0.7906



### 3.2 Path-coefficient Analysis

#### 3.2.1 Phenotypic path coefficients

The direct and indirect effect of different characters on grain yield/plant computed by using phenotypic correlations are presented in Table 3. The highest positive direct effect on grain yield per plant was exhibited by GPP (0.485), DFF(0.414), PH at the time of 45DAT (0.255), Chl (0.255), EBT (0.173), PH at the time of 30DAS (0.042), NOT (0.029), PL(0.024) the direct effects of remaining characters were too low to be considered important. GPP exhibited indirect positive effect on grain yield per plant via PH at the time of maturity (0.161), Chl (0.064), EBT (0.02), DTFI (0.002), PH at the time of 15 DAS(0.002), PH at the time of 30 DAS(0.001), SPP(0.001), High direct effect of filled spikelets/panicle on single plant yield was reported by Eidi Kohnaki et al. [13], Kiani and Nematzadeh [14], Seyoum et al. [15], Bagheri et al. [16], Bhadru et al. [17] and Chandra et al. [18].

#### 3.2.2 Genotypic path coefficients

The highest positive direct effect on grain yield per plant was exhibited by DTFI (1.034), GPP(0.372), SPP(0.358), PL(0.295), PH at the time of maturity(0.187), EBT(0.135), PH at the time of 30DAS (0.100), Chl (0.046). DTFI exhibited indirect positive effect on grain yield per plant via EBT, PH at the time of maturity, NOT, SPP, PL, PH at the time of 30 DAS, similar result supported that Bhadru et al. [17] and Chandra et al. [18] reported positive direct effect of days to 50% flowering and Eidi kohnaki et al. [13] and Nematzadeh (2012) found the positive direct effect.

The estimated residual effect was 0.790 indicating that about 80% of the variability in grain yield was contributed by the characters studied in path analysis. This residual effect towards yield in the present study might be due to many reasons, such as other characters, which are not included in the investigation, environmental factor and sampling errors. Within the scope of path analysis carried out in the present study, it is therefore, suggested that number of spikelets per panicle (SPP) and number of tillers (NOT), the main components of grain yield should be given high priority in the selection programme.

### 4. CONCLUSION

At genotypic and phenotypic level, number of grain per panicle, chlorophyll content, ear bearing tillers per plant showed highly positive significant correlation with yield per plant. Path analysis identified the highest positive direct effect on grain yield per plant with number of grain per panicle, days to 50% flowering, plant height, chlorophyll content, ear bearing tillers per plant, plant height at 30DAS, number of tillers per plant, panicle length.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Khush GS and Virk PS. Rice breeding achievements and future strategies. *Crop Improvement*. 2000;27(2):115-144.
2. Singh R, Yadav V, Mishra DN and Yadav A. Correlation and Path Analysis Studies in Rice (*Oryza sativa* L.). *Journal of Pharmacognosy and Phytochemistry*. SP1. 2018;2084-2090.
3. Singh RK and Chaudhary BD. Biometrical methods in quantitative genetic analysis. Kalyani Publishers New Delhi. 1995;215-218.
4. Dewey DR, Lu KHA. Correlation and path coefficient analysis of components of crested wheat grass seed production. *Agron. J.* 1959;51:515-518.
5. Borbora TK, Hazarika GN, Medhi AK. Correlation and path analysis for panicle characters in rice. *Crop Res.* 2005;30(2): 215-222.
6. Dhurai SY, Reddy DM and Ravi S. Correlation and Path Analysis for Yield and Quality Characters in Rice (*Oryza sativa* L.). *Rice Genom. Genet.* 2016; 7(4):1-6.
7. Namita P, Koutu GK and Akanksha T. Genetic variability, correlation and path coefficient studied for grain yield and other yield attributing traits in rice (*Oryza sativa* L.) *Int. J. Agric. Sci.* 2016;8(55): 2988-2992.
8. Nikhil B, Rangare N, and Saidaiah P. Correlation and path analysis in rice (*Oryza sativa* L.). *Int. J. Tropical Agric.* 2014;32:1-5.

9. Sathisha T. Association and path coefficient analysis for nutritional quality, grain yield and its attributing traits in traditional land races of rice. *Int. J. Agric. Sci.* 2015;7(13):841-847.
10. Qamar-Zia U, Cheema AA, Ashraf M, Rashid M, Tahir GR. Association analysis of some yield influencing traits in aromatic and non-aromatic rice. *Pak. J Bot.* 2005; 37(3):613-627.
11. Ramakrishnan S, Ananda H, Kumar CR, Sarvanan S, Malini N. Association analysis of some yield traits in Rice (*Oryza sativa* L.). *Journal of Applied Sciences Research.* 2006;2(7):402-404.
12. Mahto RN, Yadava MS, Mohan KS. Genetic variation, character association and path analysis in rainfed upland rice. *Indian J Dryland Agric. Res. and Devel.* 2003;18(2):196-198.
13. Eidi Kohnaki M, Kiani G and Nematzadeh G. Relationship between Morphological Traits in Rice Restorer Lines at F3 Generation using Multivariate Analysis. *Int J Adv Biol Biom Res.* 2013;1(6):572-577.
14. Kiani G and Nematzadeh G. Correlation and Path Coefficient Studies in F Populations of Rice. *Not Sci Biol.* 2012; 4(2):124-127.
15. Seyoum M, Alamerew S and Bantte K. Genetic Variability, Heritability, Correlation Coefficient and Path Analysis for Yield and Yield Related Traits in Upland Rice (*Oryza sativa* L.). *Journal of Plant Sciences.* 2012;7(1): 13-22.
16. Bagheri N, Babaeian-Jelodar N, Pasha A. Path coefficient analysis for yield and yield components in diverse rice (*Oryza sativa* L.) genotypes. *Bih Biol.* 2011;5:32-35.
17. Bhadru D, Reddy DL and Ramesha MS. Correlation and path coefficient analysis of yield and yield contributing traits in rice hybrids and their parental lines. *Electronic Journal of Plant Breeding.* 2011;2(1):112-116.
18. Chandra BS, Reddy TD, Ansari NA and Kumar SS. Correlation and path coefficient analysis for yield and yield components in rice (*Oryza sativa* L.). *Agric. Sci. Digest.* 2009;29(1):45-47.

© 2022 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/88495>